

Ausgabe: ZWP 10/10

Thema: Parodontalen Gewebeabbau rechtzeitig erkennen

Autor: Dr. Hans Sellmann

Literatur

Hakkarainen K et al. Collagenase activity and protein content of sulcular fluid after scaling and occlusal adjustment of teeth with deep periodontal pockets. *J Periodontal Res.* 1988 May;23(3):204-10.

Ingman T et al. Matrix metalloproteinases and their inhibitors in gingival crevicular fluid and saliva of periodontitis patients. *J Clin Periodontol.* 1996 Dec;23(12):1127-32.

Lang NP et al. Absence of bleeding on probing. An indicator of periodontal stability. *J Clin Periodontol.* 1990 Nov;17(10):714.

Mäntylä P et al. Gingival crevicular fluid collagenase-2 (MMP-8) test stick for chair-side monitoring of periodontitis. *J Periodontal Res.* 2003 Aug;38(4):436-9.

Mäntylä P et al. Monitoring periodontal disease status in smokers and nonsmokers using a gingival crevicular fluid matrix metalloproteinase-8-specific chair-side test. *J Periodontal Res.* 2006 Dec;41(6):503-12.

Netuschil L. Parodontitis- und Periimplantitis-Frühwarnung. Welcher diagnostischer Test bringt welche Aussage? *Plaque&Care* 2009: 5-9.

Reich E, Arweiler N, Benz C, Lynch E, Scholz V. Stellenwert des Chlorhexidins in der Implantatversorgung. Konsensuspapier. *ZWR* 2007; 116 (7-8): 359-362.

Sorsa T et al. Scientific basis of a matrix metalloproteinase-8 specific chair-side test for monitoring periodontal and peri-implant health and disease. *Ann N Y Acad Sci.* 1999 Jun 30;878:130-40.

Sorsa T et al. Matrix metalloproteinases (MMPs) in oral diseases. *Oral Dis.* 2004 Nov;10(6):311-8.

Sorsa T et al. Detection of gingival crevicular fluid MMP-8 levels with different laboratory and chair-side methods. *Oral Dis.* 2010 Jan;16(1):39-45.