

## **Mikroinvasive Infiltrationstherapie einer initialen Karies**

**Prof. Dr. Andrej M. Kielbassa**

### **Literaturliste**

1. Croll TP: Bonded resin sealant for smooth surface enamel defects: new concepts in "microrestorative" dentistry. *Quintessence Int*, 1987; 18: 5-10.
2. Davila JM, Buonocore MG, Greeley CB, Provenza DV: Adhesive penetration in human artificial and natural white spots. *J Dent Res*, 1975; 54: 999-1008.
3. Espelid I, Tveit AB: Clinical and radiographic assessment of approximal carious lesions. *Acta Odontol Scand*, 1986; 44: 31-37.
4. Garcia-Godoy F, Summitt JB, Donly KJ: Caries progression of white spot lesions sealed with an unfilled resin. *J Clin Pediatr Dent*, 1997; 21: 141-143.
5. Kielbassa AM, Muller J, Gernhardt CR: Closing the gap between oral hygiene and minimally invasive dentistry: A review on the resin infiltration technique of incipient (proximal) enamel lesions. *Quintessence Int*, 2009; 40: 663-681.
6. Kielbassa AM, Paris S, Lussi A, Meyer-Lueckel H: Evaluation of cavitations in proximal caries lesions at various magnification levels in vitro. *J Dent*, 2006; 34: 817-822.
7. Klimek J. Initiale Karies und ihre Behandlung. In Heidemann D (ed.). Deutscher Zahnärztekalender 1997, Hanser, München 1997: S. 23-42.
8. Martignon S, Ekstrand KR, Ellwood R: Efficacy of sealing proximal early active lesions: an 18-month clinical study evaluated by conventional and subtraction radiography. *Caries Res*, 2006; 40: 382-388.
9. Mejare I, Kallestal C, Stenlund H, Johansson H: Caries development from 11 to 22 years of age: a prospective radiographic study. Prevalence and distribution. *Caries Res*, 1998; 32: 10-16
10. Meyer-Lueckel H, Paris S, Kielbassa AM: Surface layer erosion of natural caries lesions with phosphoric and hydrochloric acid gels in preparation for resin infiltration. *Caries Res*, 2007; 41: 223-230.
11. Meyer-Lueckel H, Paris S, Mueller J, Colfen H, Kielbassa AM: Influence of the application time on the penetration of different dental adhesives and a fissure sealant into artificial subsurface lesions in bovine enamel. *Dent Mater*, 2006; 22: 22-28.
12. Mueller J, Meyer-Lueckel H, Paris S, Hopfenmuller W, Kielbassa AM: Inhibition of lesion progression by the penetration of resins in vitro: influence of the application procedure. *Oper Dent*, 2006; 31: 338-345.

13. Paris S, Meyer-Lueckel H, Colfen H, Kielbassa AM: Penetration coefficients of commercially available and experimental composites intended to infiltrate enamel carious lesions. *Dent Mater*, 2007; 23: 742-748.
14. Paris S, Meyer-Lueckel H, Kielbassa AM: Resin infiltration of natural caries lesions. *J Dent Res*, 2007; 86: 662-666.
15. Paris S, Meyer-Lueckel H, Mueller J, Hummel M, Kielbassa AM: Progression of sealed initial bovine enamel lesions under demineralizing conditions in vitro. *Caries Res*, 2006; 40: 124-129.
16. Robinson C, Brookes SJ, Kirkham J, Wood SR, Shore RC: In vitro studies of the penetration of adhesive resins into artificial caries-like lesions. *Caries Res*, 2001; 35: 136-141.
17. Robinson C, Hallsworth AS, Weatherell JA, Kunzel W: Arrest and control of carious lesions: a study based on preliminary experiments with resorcinol-formaldehyde resin. *J Dent Res*, 1976; 55: 812-818.
18. Schmidlin PR, Zehnder M, Zimmermann MA, Zimmermann J, Roos M, Roulet JF: Sealing smooth enamel surfaces with a newly devised adhesive patch: a radiochemical in vitro analysis. *Dent Mater*, 2005; 21: 545-550.
19. Waggoner WF, Ashton JJ: Predictability of cavitation based upon radiographic appearance: comparison of two film types. *Quintessence Int*, 1989; 20: 55-60.