

Kieferaugmentationen mit allogenem Knochenblöcken

Ein alternativer Ansatz zum bisherigen Goldstandard

Dr. Phillip Wallowy, Dr. Dr. Andreas Dorow/Waldshut

Implantologie Journal 4/2010

Literatur

1. Boyne PJ. The use of marrow-cancellous grafts in the regeneration of mandibular bone. *Trans Int Conf Oral Surg.* 1973;4:58-63.
2. Barone A, Varanini P, Orlando B, Tonelli P, Covani U. Deep-frozen allogeneic onlay bone grafts for reconstruction of atrophic maxillary alveolar ridges: a preliminary study. *J Oral Maxillofac Surg.* Jun 2009;67(6):1300-1306.
3. Sasso RC, LeHuec JC, Shaffrey C. Iliac crest bone graft donor site pain after anterior lumbar interbody fusion: a prospective patient satisfaction outcome assessment. *J Spinal Disord Tech.* Feb 2005;18 Suppl:S77-81.
4. Silber JS, Anderson DG, Daffner SD, et al. Donor site morbidity after anterior iliac crest bone harvest for single-level anterior cervical discectomy and fusion. *Spine (Phila Pa 1976).* Jan 15 2003;28(2):134-139.
5. Skaggs DL, Samuelson MA, Hale JM, Kay RM, Tolo VT. Complications of posterior iliac crest bone grafting in spine surgery in children. *Spine (Phila Pa 1976).* Sep 15 2000;25(18):2400-2402.
6. Albert A, Leemrijse T, Druez V, Delloye C, Cornu O. Are bone autografts still necessary in 2006? A three-year retrospective study of bone grafting. *Acta Orthop Belg.* Dec 2006;72(6):734-740.
7. Gomes KU, Carlini JL, Biron C, Rapoport A, Dedivitis RA. Use of allogeneic bone graft in maxillary reconstruction for installation of dental implants. *J Oral Maxillofac Surg.* Nov 2008;66(11):2335-2338.
8. U.Schmid, F.Thielemann, P.Sprung, et al. Multizentrische prospektive Sammelstudie über die Knochenrekonstruktion mit vitalisierten Allografts: klinische Ergebnisse und ökonomische Aspekte. *68 Kongress der Unfallchirurgie.* 2004.
9. Putzier M, Strube P, Funk JF, et al. Allogenic versus autologous cancellous bone in lumbar segmental spondylodesis: a randomized prospective study. *Eur Spine J.* May 2009;18(5):687-695.
10. Haynert W. [Application possibilities of human bone-collagen substance as a transplant in skeletal surgery]. *Beitr Orthop Traumatol.* Aug 1990;37(8):453-461.

11. Kuhls R, Werner-Rustner M, Kuchler I, Soost F. Human demineralised bone matrix as a bone substitute for reconstruction of cystic defects of the lower jaw. *Cell Tissue Bank.* 2001;2(3):143-153.
12. Thielicke U, Thielicke B, von Versen R, Denner K. [Clinical study on the application of demineralized bone matrix (DBM) in surgical orthodontics]. *Beitr Orthop Traumatol.* Aug 1990;37(8):461-465.
13. Wildemann B, Kadow-Romacker A, Pruss A, Haas NP, Schmidmaier G. Quantification of growth factors in allogenic bone grafts extracted with three different methods. *Cell Tissue Bank.* 2007;8(2):107-114.
14. Pruss A, Perka C, Degenhardt P, et al. Clinical efficacy and compatibility of allogeneic avital tissue transplants sterilized with a peracetic acid/ethanol mixture. *Cell Tissue Bank.* 2002;3(4):235-243.
15. Soost, Frank. Validierung des Knochenumbaus von Knochenersatzmaterialien in der Mund-, Kiefer- und Gesichtschirurgie. Habilitationsschrift, 2006, Humboldt-Universität, Berlin.
16. Simion M, Fontana F, Maiorana C. Vertical ridge augmentation by expanded-polytetrafluoroethylene membrane and a combination of intraoral autogenous bone graft and deproteinized anorganic bovine bone (Bio Oss). *Clin Oral Implants Res.* 2007 oct; 18(5): 620-9.
17. Meyer B: Reinigungsleistung Peressigsäure basierter Desinfektionsmittel für die Aufbereitung medizinischer Instrumente. *Hyg Med* 2004; 29: 106–112