

## **Laser zur Kariesdetektion: für okklusale und proximale Läsionen geeignet?**

***Dr. med. dent. Jonas Almeida Rodrigues, Prof. Adrian Lussi, Dr. Klaus Neuhaus/Bern, Schweiz, Dr. Michele Baffi Diniz/São Paulo, Brasilien***

***Laser Journal 3/2010***

### Bibliographie

1. Aljehani A, Tranaeus S, Forsberg CM, Angmar-Mansson B, Shi XQ: In vitro quantification of white spot enamel lesions adjacent to fixed orthodontic appliances using quantitative light-induced fluorescence and DIAGNOdent. Acta Odontol Scand 62: 313-318 (2004).
2. Alkurt M, Peker I, Deniz Arisu H, Bala O, Altunkaynak B: In vivo comparison of laser fluorescence measurements with conventional methods for occlusal caries detection. Lasers Med Sci 23: 307-312 (2008).
3. Bader JD, Shugars DA: A systematic review of the performance of a laser fluorescence device for detecting caries. J Am Dent Assoc: 135: 1413-1426 (2004)
4. Bamzahim M, Aljehani A, Shi XQ: Clinical performance of DIAGNodent in the detection of secondary carious lesions. Acta Odontol Scand 63: 26-30 (2005).
5. Bamzahim M, Shi XQ, Angmar-Mansson B: Secondary caries detection by DIAGNOdent and radiography: a comparative in vitro study. Acta Odontol Scand 62: 61-64 (2004).
6. Boston DW: Initial in vitro evaluation of DIAGNOdent for detecting secondary carious lesions associated with resin composite restorations. Quintessence Int 34: 109-116 (2003).
7. Braga MM, Mendes FM, Martins CR, Imparato JC: Effect of the calibration method of a laser fluorescence device for detecting occlusal caries in primary molars. Pediatr Dent 28: 451-454 (2006).
8. Braun A, Krause F, Jepsen S: The Influence of the Calibration Mode of a Laser Fluorescence Device on Caries Detection. Caries Res 39: 144-149. (2005).
9. Diniz MB, Rodrigues JA, Hug I, Cordeiro RC, Lussi A: The influence of pit and fissure sealants on infrared fluorescence measurements. Caries Res 42: 328-333 (2008).
10. Diniz MB, Rodrigues JA, Paula AB, Cordeiro RD: In vivo evaluation of laser fluorescence performance using different cut-off limits for occlusal caries detection. Lasers Med Sci 24: 295-300 (2009).
11. Hibst R, Paulus R, Lussi A: A detection of occlusal caries by laser fluorescence: basic and clinical investigations. Med Laser Appl 16: 295-213 (2001).
12. Jablonski-Momeni A, Stachniss V, Ricketts DN, Heinzl-Gutenbrunner M, Pieper K: Reproducibility and accuracy of the ICDAS-II for detection of occlusal Caries in vitro. Caries Res 42: 79-87 (2008).

13. Kuhnisch J, Bucher K, Henschel V, Hickel R: Reproducibility of DIAGNOdent 2095 and DIAGNOdent pen measurements: results from an in vitro study on occlusal sites. *Eur J Oral Sci* 115: 206-211 (2007).
14. Kuhnisch J, Bucher K, Hickel R: The intra/inter-examiner reproducibility of the new DIAGNOdent Pen on occlusal sites. *J Dent* 35: 509-512 (2007).
15. Lennon ÁM, Buchalla W, Switalski L, Stookey GK: Residual caries detection using visible fluorescence. *Caries Res* 36: 315-319 (2002).
16. Lussi A, Francescut P: Performance of conventional and new methods for the detection of occlusal caries in deciduous teeth. *Caries Res* 37: 2-7 (2003).
17. Lussi A, Hack A, Hug I, Heckenberger H, Megert B, Stich H: Detection of approximal caries with a new laser fluorescence device. *Caries Res* 40: 97-103 (2006).
18. Lussi A, Hellwig E: Performance of a new laser fluorescence device for the detection of occlusal caries in vitro. *J Dent* 34: 467-471 (2006).
19. Lussi A, Imwinkelried S, Pitts N, Longbottom C, Reich E: Performance and reproducibility of a laser fluorescence system for detection of occlusal caries in vitro. *Caries Res* 33: 261-266 (1999).
20. Neuhaus K, Longbottom C, Ellwood R, Lussi A: Novel lesion detection aids. *Monogr Oral Sci* 21: 52-62 (2009).
21. Novaes TF, Matos R, Braga MM, Imparato JCP, Raggio DP, Mendes FM: Performance of a pen-type laser fluorescence device and conventional methods in detecting approximal caries lesions in primary teeth -- in vivo study. *Caries Res* 43: 36-42 (2009).
22. Pinelli C, Campos Serra M, de Castro Monteiro Loffredo L: Validity and reproducibility of a laser fluorescence system for detecting the activity of white-spot lesions on free smooth surfaces in vivo. *Caries Res* 36: 19-24 (2002).
23. Rodrigues JA, Diniz MB, Hug I, Cordeiro RCL, Lussi A: Relationship between DIAGNOdent values and sealant penetration depth on occlusal fissures. *Clinical Oral Investigations* DOI 10.1007/s00784-009-0321-1 (2009).
24. Rodrigues JA, Hug I, Diniz MB, Cordeiro RCL, Lussi A: The influence of zero-value subtraction on the performance of two laser fluorescence devices for detecting occlusal caries in vitro. *J Am Dent Assoc* 139: 1105-1112 (2008).
25. Rodrigues JA, Hug I, Diniz MB, Lussi A: Performance of fluorescence methods, radiographic examination and ICDAS II on occlusal surfaces in vitro. *Caries Res* 42: 297-304 (2008).
26. Shi XQ, Tranaeus S, Angmar-Mansson B: Comparison of QLF and DIAGNOdent for quantification of smooth surface caries. *Caries Res* 35: 21-26 (2001)
27. Shi XQ, Tranaeus S, Angmar-Mansson B: Validation of DIAGNOdent for quantification of smooth-surface caries: an in vitro study. *Acta Odontol Scand* 59: 74-78 (2001).

28. Wicht MJ, Haak R, Stutzer H, Strohe D, Noack MJ: Intra- and interexaminer variability and validity of laser fluorescence and electrical resistance readings on root surface lesions. *Caries Res* 36: 241-248 (2002).