

Über den Tellerrand blacken

Fraktionierte CO₂-Lasertherapie bei lichtgeschädigter Haut und Falten

Dr. med. Kai Rezai/Münster

Laser Journal 2/2010

Literaturverzeichnis

- 1 Maimann T.H.: Stimulated Optical Radiation in Ruby; Natur (1960) 187 (4736) 493–494
- 2 Maiman T.H.: The Laser Odyssey. Laser Press 2001
- 3 Goldman L., D. J. Blaney, D. J. Kindel jun., D. Richfield & E. K. Franke: Pathology of the Effect of the Laser Beam on the Skin; Nature (1963) 197, 912–914
- 4 Alster TS, West TB. Resurfacing of atrophic facial acne scars with a highenergy,pulsed carbon dioxide laser. Dermatol. Surg. 1996; 22: 151–154
- 5 Waldorf HA, Kauvar AN, Geronemus RG. Skin resurfacing of fine to deep rhytides using a char-free carbon dioxide laser in 47 patients. Dermatol. Surg. 1995; 21: 940–946
- 6 Kuo T, Speyer MT, Ries WR, Reinisch L. Collagen thermal damage and collagen synthesis after cutaneous laser resurfacing. Lasers Surg. Med. 1998; 23: 66–71
- 7 Smith KJ, Skelton HG, Graham JS, Hurst CG, Hackley BE, Jr. Increased smooth muscle actin, factor XIIIa, and vimentin-positive cells in the papillary dermis of carbon dioxide laser-debrided porcine skin. Dermatol. Surg. 1997; 23: 891–895
- 8 Khatri KA, Ross V, Grevelink JM, Magro CM, Anderson RR. Comparison of erbium:YAG and carbon dioxide lasers in resurfacing of facial rhytides. Arch. Dermatol. 1999; 135: 391–397
- 9 Lowe NJ, Lask G, Griffin ME. Laser skin resurfacing. Pre- and posttreatment guidelines. Dermatol. Surg. 1995; 21: 1017–1019
- 10 Grund A. Hochfrequente Radiomikrochirurgie in der kosmetischen Gesichtschirurgie, Face 2007 Issue1: 44–47
- 11 Manstein D, Herron GS, Sink RK, Tanner H, Anderson RR. Fractional photothermolysis: a new concept for cutaneous remodeling using microscopic patterns of thermal injury. Lasers Surg. Med. 2004; 34: 426–438
- 12 Tannous Z. Fractional resurfacing. Clin. Dermatol. 2007; 25: 480–486
- 13 Lupton JR, Williams CM, Alster TS. Nonablative laser

- 14 Bedi VP, Chan KF, Sink RK, Hantash BM, Herron GS, Rahman Z, Struck SK, Zachary CB. The effects of pulse energy variations on the dimensions of microscopic thermal treatment zones in nonablative fractional resurfacing. *Lasers Surg. Med.* 2007; 39: 145– 155
- 15 Karsai S, Raulin C. *Hautarzt* 2008 59: 92–100
- 16 Bodendorf MO, Grunewald S, Wetzig T, Simon JC, Paasch U. Fraktionale Lasertherapie der Haut. *JDDG* 2009; 7:301–308
- 17 Goerge T, Peukert N, Bayer H, Rütter A. Ablative fractional photothermolysis- A novel Stepp in skin resurfacing. *J. med. Laser App* 2008; 23 93–98
- 18 Hasegawa T, Matsukura T, Mizuno Y, Suga Y, Ogawa H, Ikeda S. Clinical trial of a laser device called fractional photothermolysis system for acne scars. *J. Dermatol.* 2006; 33: 623–627.
- 10 Grund A. Hochfrequente Radiomikrochirurgie in der kosmetischen Gesichtschirurgie, *Face* 2007 Issue1: 44–47
- 11 Manstein D, Herron GS, Sink RK, Tanner H, Anderson RR. Fractional photothermolysis: a new concept for cutaneous remodeling using microscopic patterns of thermal injury. *Lasers Surg. Med.* 2004; 34:426–438
- 12 Tannous Z. Fractional resurfacing. *Clin. Dermatol.* 2007; 25:480–486
- 13 Lupton JR, Williams CM, Alster TS. Nonablative laser