

Durchtrittsprofile korrekt einstellen

Einfluss der Provisorien auf den Erfolg der definitiven Versorgung

Dr. Orcan Yüksel/Frankfurt, Dr. Kris Chmielewski/Danzig, Polen

Implantologie Journal 8/2010

Literatur:

1. **Abrahamsson, I, Berglundh, T, and Lindhe, J:** The mucosal barrier following abutment dis/reconnection. An experimental study in dogs. *J Clin Periodontol*; 1997, 24 (8): 568-572
2. **Degidi, M, Nardi, D, and Piattelli, A:** Immediate versus one-stage restoration of small-diameter implants for a single missing maxillary lateral incisor: a 3-year randomized clinical trial. *J Periodontol*; 2009, 80 (9): 1393-1398
3. **Degidi, M, Piattelli, A, Gehrke, P, et al.:** Five-year outcome of 111 immediate nonfunctional single restorations. *J Oral Implantol*; 2006, 32 (6): 277-285
4. **Degidi, M, Scarano, A, Piattelli, M, et al.:** Bone remodeling in immediately loaded and unloaded titanium dental implants: a histologic and histomorphometric study in humans. *J Oral Implantol*; 2005, 31 (1): 18-24
5. **Zechner, W, Kneissel, M, Kim, S, et al.:** Histomorphometrical and clinical comparison of submerged and nonsubmerged implants subjected to experimental peri-implantitis in dogs. *Clin Oral Implants Res*; 2004, 15 (1): 23-33