

Literaturliste

Zwei Stufen, viele Vorteile

GC Germany/Bad Homburg

Dental Tribune German Edition 7+8/2014

1. Vergleich basiert auf Berechnungen von K. H. Friedl unter Berücksichtigung der Daten von Micheelis W., Meyer V. P. (2002): Arbeitswissenschaftliche Beanspruchungsmuster zahnärztlicher Dienstleistungen (BAZ-II) – Materialien zur Beanspruchungsdauer und Beanspruchungshöhe ausgewählter Behandlungsanlässe, IDZ-Materialeinreihe, Band 27, Deutscher Zahnärzte Verlag DÄV.
2. Diem VTK et al. (2013): The effect of a nano-filled resin coating on the 3-year clinical performance of a conventional high-viscosity glass-ionomer cement. Clinical Oral Investigations DOI 10.1007/s00784-013-1026-z.
3. Klinke T, Daboul A, Biffar R: EQUIA - RCT in the field (2013): Longevity after 24 months. CED IADR Florenz, 2013, Abstract 3. Abruf am 30.01.2014 unter: <https://iadr.confex.com/iadr/ced13/webprogram/Paper179792.html>.
4. Basso M, et al. (2013). Permanent restorations with glassionomer cements. Clinical evaluation on 319 cases, IADR Seattle, 2013, Abstract 594. Abruf am 30.01.2014 unter: <https://iadr.confex.com/iadr/13iags/webprogram/Paper175751.html>
5. Gurgan S (2013): EQUIA study – Clinical performance of EQUIA restorative System in restoration of posterior teeth. Vortrag auf CED-IADR-Meeting in Florenz: 06.09.2013.
6. Lohbauer U, Krämer N, Siedschlag G, Schubert EW, Laufer B, Müller FA, Petschelt A, Ebert J. Strength and wear resistance of a dental glass-ionomer cement with a novel nanofilled resin coating. Am J Dent. 2011; 14:124-128.
7. Diem VTK et al. (2013): The effect of a nano-filled resin coating on the 3-year clinical performance of a conventional high-viscosity glass-ionomer cement. Clinical Oral Investigations DOI 10.1007/s00784-013-1026-z.