

Ausgabe: **Cosmetic Dentistry 2/2017**

Thema: Versorgung mit Feldspatkeramikveneers bei parodontalem Attachmentverlust

Autoren: Dr. Sven Egger, M.Sc., M.Sc., ZTM Christian Berg

Literatur

1. Magne P, Belser U: Adhäsiv befestigte Keramikrestaurationen. Berlin: Quintessenz, 2004.
2. Fradeani M: Evaluation of dentolabial parameters as part of a comprehensive esthetic analysis. Eur J Esthet Dent. 2006; 1: 62-69.
3. D'Amico A: The canine teeth-normal functional relation of the natural teeth of man. J South Calif Dent Assoc. 1958; 26: 6-23, 49-60,127-142,175-182, 194-208,239-241.
4. Magne P, Magne M, Belser U: The esthetic width in fixed prothodontics. J Prosthodont 1999;8: 106-118.
5. Witkowski S, Schicha K: Prep Veneers & Non-Prep Veneers. Berlin: Quintessenz, 2010.
6. Vailati F, Belser UC: Palatal and facial veneers to treat severe dental erosion: a case report following the three-step technique and the sandwich approach. Eur J Esthet Dent 2011 Autumn;6(3):268-78.
7. Peumans M, Van Meerbeek B, Lambrechts P, Vanherle G, Quirynen M: The influence of direct composite additions for the correction of tooth form and/or position on periodontal health: A retrospective stud. J Periodontol 1998;69 :422-427.
8. Rufenacht CR: Fundamentals of Esthetics. Chicago: Quintessenz, 1990: 67-134.