

Ausgabe: ZWP Zahnarzt Wirtschaft Praxis 7+8/2014

Thema: Implantatgetragene Prothese zur Versorgung zahnloser Kiefer

Autoren: Dr. Dr. Manfred Wolf, Dr. Daniel Rein

Literatur

[1]. Attard NJ, Zarb GA. Long-term treatment outcomes in edentulous patients with implant-fixed prostheses: the Toronto study. *Int J Prosthodont* 2004;17:417-424.

[2]. Tallgren A. The continuing reduction of the residual alveolar ridges in complete denture wearers: a mixed-longitudinal study covering 25 years. *J Prosthet Dent* 1972;27:120-132.

[3]. Assuncao WG, Barao VA, Delben JA, Gomes EA, Tabata LF. A comparison of patient satisfaction between treatment with conventional complete dentures and overdentures in the elderly: a literature review. *Gerodontology* 2010;27:154-162.

[4]. Wismeijer D, Van Waas MA, Vermeeren JI, Mulder J, Kalk W. Patient satisfaction with implant-supported mandibular overdentures. A comparison of three treatment strategies with ITI-dental implants.

[5]. Wolfart M, Weber V. Abnehmbarer Zahnersatz auf Implantaten - implantatgestützte Versorgungsformen im zahnlosen Kiefer. *Logo* 2013:15-21.

International journal of oral and maxillofacial surgery 1997;26:263-267.

[6]. Thomason JM, Feine J, Exley C, Moynihan P, Muller F, Naert I, et al. Mandibular two implant-supported overdentures as the first choice standard of care for edentulous patients--the York Consensus Statement. *Br Dent J* 2009;207:185-186.