

**Ausgabe:** KN 6-2017, S. 7ff.

**Thema:** Extraktion zweiter Prämolaren – Behandlung mit selbstligierenden Brackets

**Autoren:** Dr. Dirk Kujat

---

## Literatur

1. A multi-center randomized controlled trial to compare a self-ligating bracket with a conventional bracket in a UK population: *Part I: Treatment efficiency* Lian O'Dywer, Simon J. Littlewood, Shahla Rahman, R. James Spencer, Sophy K. Barber and Joanne S. Russell *Angle Orthod.* 2016; 86:142-148
2. Treatment time, outcome, and anchorage loss comparisons of self-ligating and conventional brackets  
Ferdinand M. Machibya, Xingfu Bao, Lihua Zhao and Min Hu, *Angle Orthod.* 2013; 83: 280-285
3. Canine retraction rate with self-ligating brackets vs conventional edgewise brackets  
S. Jack Burrow, *Angle Orthod.* 2010; 80: 626-633
4. Dibbets and van der Welle *AJO* – 1991
5. Nance HN The removal of second premolars in orthodontic treatment *Am J Orthod.* 1949; 35:685-96
6. Schoppe RJ. An analysis of second premolar extraction procedures *Angle Orthod.* 1964;34:292-302
7. Salzmann JA. The rationale of extraction as an adjunct to orthodontic mechanotherapy and the sequelae of extraction in the absence of orthodontic guidance *Am J Orthod Oral Surg.* 1945;31:181-202