

Röntgen up to date: Analog oder digital?

Priv.-Doz. Dr. Andreas Bindl, Dr. Daniel Wolf/Zürich, Schweiz

Endodontie Journal 4/2010

Literatur

Silva MA, Wolf U, Heinicke F, Bumann A, Visser H, Hirsch E.
Cone-beam computed tomography for routine orthodontic treatment planning: a radiation dose evaluation.
Am J Orthod Dentofacial Orthop. 2008 May;133(5):640.e1-5.

Wörtche R, Hassfeld S, Lux CJ, Müssig E, Hensley FW, Krempien R, Hofele C.
Clinical application of cone beam digital volume tomography in children with cleft lip and palate.
Dentomaxillofac Radiol. 2006 Mar;35(2):88-94.

Bender IB, Seltzer S
Roentgenographic and direct observation of experimental lesions in bone (part I)
J Am Dent Assoc. 1961; 62:152-160

Bender IB, Seltzer S
Roentgenographic and direct observation of experimental lesions in bone (part II)
J Am Dent Assoc. 1961; 62:708-716

Lee S-J, Messer HH.
Radiographic appearance of artificially prepared periapical lesions confined to cancellous bone.
International Endodontic Journal 1986. Volume 19 Issue 2, Pages 64 - 72

Barbat J, Messer HH. Detectability of artificial periapical lesions using direct digital and conventional radiography.
J Endod. 1998 Dec;24(12):837-42.

Low KM, Dula K, Bürgin W, von Arx T.
Comparison of periapical radiography and limited cone-beam tomography in posterior maxillary teeth referred for apical surgery.
J Endod. 2008 May;34(5):557-62.

Estrela C, Bueno MR, Leles CR, Azevedo B, Azevedo JR.
Accuracy of cone beam computed tomography and panoramic and periapical radiography for detection of apical periodontitis.
J Endod. 2008 Mar;34(3):273-9.

Lofthag-Hansen S, Huumonen S, Gröndahl K, Gröndahl HG.
Limited cone-beam CT and intraoral radiography for the diagnosis of periapical pathology.
Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2007 Jan;103(1):114-9.

Patel S. New dimensions in endodontic imaging: Part 2. Cone beam computed tomography.
Int Endod J. 2009 Jun;42(6):463-75. Epub 2009 Mar 2.

Nair PN, Sjögren U, Figdor D, Sundqvist G.
Persistent periapical radiolucencies of root-filled human teeth, failed endodontic treatments, and periapical scars.
Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 1999 May;87(5):617-27.