

**Ausgabe:** face & body 3/2017

**Thema:** Der Einsatz von Fillern in der rekonstruktiven Lippenchirurgie

**Autor:** Prof. Dr. Dr. Niels Christian Pausch

---

**Literatur:**

1. Feistl W. Verbesserung der Lippenästhetik bei Patienten mit Lippen-Kiefer-Gaumenspalten durch Permanent Make-up und Lippenaugmentation mittels Hyaluronsäure-Filler. *Face* 4/2012: 16-20
2. Niechajev I. Lip enhancement. Surgical alternatives and histologic aspects. *Plast Reconstr Surg.* 2000, 105: 1173–83
3. Shab A. Lippenaugmentation mit der neuen Füllergeneration Agarose-Gel. *Face* 2/2016: 16-18
4. Patel IA, Hall PN. Free dermis-fat graft to correct the whistle deformity in patients with cleft lip. *Br J Plast Surg.* 2004, 57: 160–4
5. Staebel C, Verheyden CN. The use of dermal fat grafts for the correction of secondary cleft lip deformities. *Plast Reconstr Surg.* 2009, 123: 151e–2e
6. Abdali H, Kachuee AAA, Mohammadi-Mofrad R, Houghoughi MA, Esmalian-Afyouni N. Surgical correction of upper lip deficit in patients with cleft lip using dermis fat graft. *J Res Med Sci.* 2017, 22: 29
7. Jones CM, Morrow BT, Albright WB, Long RE, Samson TD, Mackay DR. Structural fat grafting to improve reconstructive outcomes in secondary cleft lip deformity. *Cleft Palate Craniofac J.* 2016, 54: 70–4