

**Ausgabe:** KN 9-2017, S. 1ff.  
**Thema:** Surgery First in der Behandlung skelettaler Dysgnathien  
**Autoren:** Priv.-Doz. Dr. Dr. Robin Seeberger

---

### Literatur

- Sharma VK, Yadav K, Tandon P: An overview of surgery-first approach: Recent advances in orthognathic surgery. *J Orthod Sci* 4: 9 12, 2015.
- Huang CS, Hsu SS, Chen YR: Systematic review of the surgery-first approach in orthognathic surgery. *Biomed J* 37:184 190, 2014.
- Bell WH, Creekmore TD: Surgical-orthodontic correction of mandibular prognathism. *Am J Orthod* 63:256 270, 1973.
- Worms FW, Isaacson RJ, Speidel TM: Surgical orthodontic treatment planning: profile analysis and mandibular surgery. *Angle Orthod* 46:1 25, 1976.
- Proffit WR, Miguel JA: The duration and sequencing of surgical-orthodontic treatment. *Int J Adult Orthodon Orthognath Surg* 10:35, 1995.
- Bohannon HM: Studies in the alteration of vestibular depth I. Complete denudation. *J Periodontol* 33:354 359, 1962.
- Hoogeveen EJ, Jansma J, Ren Y: Surgically facilitated orthodontic treatment: a systematic review. *Am J Orthod Dentofacial Orthop* 145:51 64, 2014.
- Meikle MC: The tissue, cellular, and molecular regulation of orthodontic tooth movement: 100 years after Carl Sandstedt. *Eur J Orthod* 28:221 240, 2006.
- Giannopoulou C, Dudic A, Kiliaridis S: Pain discomfort and crevicular fluid changes induced by orthodontic elastic separators in children. *J Pain* 7:367 376, 2006.
- Di Domenico M, D'Apuzzo F, Feola A, Cito L, Monsurro A, Pierantoni GM, Berrino L, De Rosa A, Polimeni A, Perillo L: Cytokines and VEGF induction in orthodontic movement in animal models. *J Biomed Biotechnol* 2012:Article ID 201689, 2012.
- Hernández-Alfaro F, Guijarro-Martínez R, Peiró-Guijarro MA : Surgery First in Orthognathic Surgery: What Have We Learned? A Comprehensive Workflow Based on 45 Consecutive Cases. *J Oral Maxillofac Surg.* 2014 Feb;72(2):376-90
- Park JK, Choi JY, Yang IH, Baek SH: Patient's Satisfaction in Skeletal Class III Cases Treated With Two-Jaw Surgery Using Orthognathic Quality of Life Questionnaire: Conventional Three-Stage Method Versus Surgery-First Approach. *J Craniofac Surg* 26:2086 2093, 2015.

Zingler S, Hakim E, Finke D, Brunner M, Saure D, Hoffmann J, Lux CJ, Erber R, Seeberger R.:

Surgery-first approach in orthognathic surgery: Psychological and biological aspects - A prospective cohort study. J Craniomaxillofac Surg. 2017 Jun 6.

Nagasaka H, Sugawara J, Kawamura H, et al: "Surgery first" skeletal Class III correction using the Skeletal Anchorage System. *J Clin Orthod* 43:97, 2009

Dysgnathie–OP's ohne vorherige KFO, M. Schüßler, R.B. Drommer, *Kieferorthopädische Nachrichten* 7, II, 2009

Seeberger R, Asi Y, Thiele OC, Hoffmann J, Stucke K, Engel M.: Neurosensory alterations and function of the temporomandibular joint after high oblique sagittal split osteotomy: an alternative technique in orthognathic surgery. *Br J Oral Maxillofac Surg*. 2013 Sep;51(6):536-40.

Kuehle R, Berger M, Saure D, Hoffmann J, Seeberger R.: High oblique sagittal split osteotomy of the mandible: assessment of the positions of the mandibular condyles after orthognathic surgery based on cone-beam tomography. *Br J Oral Maxillofac Surg*. 2016 Jul;54(6):638-42. doi: 10.1016/j.bjoms.2016.03.017. Epub 2016 Apr 3.

Hunt OT, Johnston CD, Hepper PG, Burden DJ: The psychosocial impact of orthognathic surgery: a systematic review. *Am J Orthod Dentofacial Orthop* 120:490 497, 2001.

Soh CL, Narayanan V: Quality of life assessment in patients with dentofacial deformity undergoing orthognathic surgery--a systematic review. *Int J Oral Maxillofac Surg* 42:974 980, 2013.