

Ausgabe: DENTALZEITUNG 4/17, S. 64ff.

Thema: Gesamtkonzept zur Therapie von Klasse-II-Füllungen

Autor: Dr. Christian Ehrensberger

Literatur

1. American Dental Association Procedure Recap Report (2006).
2. DentalTown (2012). Restorative Dentistry. Monthly Poll: What is the most challenging part of a Class II Restoration?
3. Usman et al, Sensitivity in composite restorations, Pakistan Oral & Dental Journal Vol 34, No. 3 (09/2014); Berkowitz et al, Postoperative Hypersensitivity in Class I Resin-based Composite Restorations, Compend Contin Educ Dent. 2009; 30(6): 356–363; Haller, Die Postoperative Hypersensibilität, zm 99, Nr. 6a, 13.03.2009 S. 44-51; Briso et al, Clinical Assessment of Postoperative Sensitivity in Posterior Composite Restorations, Operative Dentistry, 2007, 32-5, 421-426.
4. Council on Scientific Affairs of the American Dental Association. Spring 2009;4(2).
5. Dentsply Sirona Restorative, Anwenderumfrage 2015 (297 Zahnärzte in Europa).
6. JWV van Dijken, „Class I and Class II SDR/CeramX mono+ restorations bonded with the one step self-etching adhesive XenoV+, five year report“, Biomaterial Research Group, Umeå, Schweden, 15. Februar 2016. Veröffentlicht auf dem Kongress der International Association for Dental Research (IADR) 2016 in Seoul, Republik Korea.