

**Die interdisziplinäre Therapie bei der Rehabilitation der dentofazialen Ästhetik bei erwachsenen Patienten**

**Autoren**\_Prof. Dr. med. dent. Nezar Watted, Bad Mergentheim, Dr. med. dent. Tobias Teuscher, Bamberg, Prof. Dr. med. Dr. med. dent. Josip Bill, Würzburg

---

**Face 2/2010**

## **Literatur**

1. Bell, H.W., Jacobs, J.D.: Simultaneous repositioning of the maxilla, mandible, and chin. Treatment planning and analysis of the soft tissues. Am J Orthod 91, 28-50 (1986).
2. Bell, W.H., Proffit, H.R., White, R.P.: Surgical correction of facial deformities. Saunder Verlag, Baltimor (1980).
3. Bell, W.H., Schdeman, G.B.: Correction of vertical maxillary deficiency: Stability and soft tissue changes. J Oral surg 39, 666-670 (1981).
4. Buddgaard, M., Melsen, B., Terp, S.: Changes during and following total maxillary osteotomy (Le Fort I Procedure): a cephalometric study. Eur J Orthod 8, 21-29 (1986).
5. Burstone, G.J.: The integumental profile. Am J Orthod 44, 1-25 (1958).
6. Burstone, G.J.: Lip posture and its significance in treatment planning. Am J Orthod 53, 262-84 (1967).
7. Calangna, L., Silverman, S., Garfinkel, L.: Influence of neuromuscular conditioning on centric relation registrations. J Prosthet Dent 30, 598-604 (1973).
8. Carlotti, A.E., Aschaffenburg, P.H., Schendel, S.A.: Facial changes associated with surgical advancement of the lip and maxilla. J Oral Maxillofac Surg 44, 593-596 (1986).
9. Carlson D.S., Ellis, E., Dechow, P.C.: Adptation of the suprathyroid muscle complex to mandibular advancement surgery. Am J Orthod 92, 134-143 (1987).
10. Celenza, F.V.: The centric position. replacement and character. J Prosthet Dent 30, 591-598 (1973).
11. Coh-Stock, C.: Die chirurgische immediate Regulierung der Kiefer, chirurgische Behandlung der Prognathie. Dtsch Vjschr Zahnheik 37, 320-331 (1921).
12. Collins, P., Epker, B.N.: The alar basse Cinch: A technique for prevention of alar base flaring secondary to maxillary surgery. Oral Surg 53, 549-554 (1982).

13. Dal Pont G.: L`osteotomia retromolare per la correzione della progenia, Minerva chir. 18, 1138-1141 (1959).
14. Dal Pont G.: Die retromolare Osteotomie zur Korrektur der Progenie, der Retrogenie und des Mordex apertus. Öst Z Stoma 58, 8-10 (1961).
15. Dann, J.J., Fonseca, R.J., Bell, W.H.: Soft tissue changes associated with total maxillary advancement: a preliminary study. J Oral Surg 34, 19-23 (1976).
16. De Assis, E.A., Starck, W.J., Epker, B.: Cephalometric analysis of profile nasal esthetics. Part III. Postoperative changes after isolated superior repositioning. Int J Adult Orthod Orthognath 11, 279-288 (1996).
17. Ellis, E.: Condylar Positioning Devices for Orthodontic Surgery: Are They Necessary. J Oral Maxillofac Syrg 52, 536-552 (1994).
18. Ellis, E., Hinton R.J.: Histologic examination of the temporomandibular joint after mandibular advancement with and without rigid fixation: An experimental investigation in adult Macaca mulatta. J Oral Maxillofac Syrg 49, 1316 (1991).
19. Epker, B.: Esthetic maxillofacial surgery. Lea und Febiger Verlag, Philadelphia (1994).
20. Epker, B. N., Fish, L. C.: Surgical-orthodontic correction of open-bite deformity. Am J Orthod 71, 278-299 (1977).
21. Epker, B.N., Fish, L.C.: Surgical superior repositioning of the maxilla: What to do with the mandible?. Am j Orthod 78, 164-191 (1980).
22. Freihofer, H.P.M.: Changes in nasal Profile after maxillary advancement in cleft and non-cleft patients. J Maxillofac Surg 5, 20-27 (1977).
23. Friede, H., Kahnberg, K.-E., Adell, R., Riedell, A.: Accuracy of cephalometric prediction in orthognathic surgery. J Oral Maxillofac Surg 45, 754-760 (1987).
24. Hogeman, K.E., Sarnäs, K.V., Willmar, K.: One-stage surgical and dental-orthopedic correction of bimaxillary and craniofacial dysostosis. In: Fortschr Kiefer Gesichtschr, Bd 18: 39-49, Thieme Verlag, Stuttgart (1974)
25. Hui, E., Hägg, E.U.O. Tidman, H: Soft tissue changes following maxillary osteotomies in cleft lip and palate and non-cleft Patients. J Craniomaxillofac Surg 22, 182-186 (1994).
26. Jensen, A.C., Sinclair, P.M., Wolford, L.M.: Soft tissue changes associated with double jaw surgery. Am J Orthod Dentofac Orthop 101, 266-275 (1992).

27. Lee, D.Y., Bailey, L.J., Proffit, W.R.: Soft tissue changes after superior repositioning of the Maxilla with Le Fort I osteotomy: 5-years follow-up. *Int J Adult Orthod Orthognath* 11, 301-311 (1996).
28. Legan, H.L., Burstone, G.J.: Soft tissue cephalometric analysis for orthognathic surgery. *J Oral Surg*; 38, 744-51 (1980).
29. Lindorf, H.H.: Funktionsstabile Tandem-Verschraubung der sagittalen Ramusosteotomie –Operationsthenik, neue Instrumente und Erfahrungen. *Dtsch Z Mund Kiefer Gesichtschir*; 8, 367-373 (1984).
30. Lindorf, H.H.: Sagittal ramus osteotomy with tandem screw fixation –technique and results. *J. Maxillofac Surg* 14, 311-316 (1986).
31. Obwegeser, H.: The surgical correction of mandibular prognathism and retrognathia with consideration of genioplasty. *J Oral Surg* 10, 677-689 (1957).
32. Obwegeser, H.: The indication for surgical correction of mandibular deformity by sagittal splitting technique. *Br J Surg* 1, 157-168 (1963).
33. Obwegeser, H., Trauner, R.: Zur Operationstechnik bei der Progenie und anderen Unterkieferanomalien. *Dtsch Zahn Mund Kieferheilk* 23, 1-26 (1955).
34. Radney, L.J., Jacobs, J.D.: Soft tissue changes associated with surgical total maxillary intrusion. *Am J Orthodont* 80, 191-212 (1981)
35. Rekow E.D., Speidel, T.M., Koenig, R.A.: Location of the mandibular center of Autorotation in maxillary impaction surgery. *Am J Orthod Dentofac Orthop* 103, 530-536 (1993).
36. Rosen, H.M.: Lip-nasal esthetics following Le Fort I Osteotomy. *Plast Reconstr Surg* 81, 171-182 (1988)
37. Schuchardt, K.: Formen des offenen Bisses und ihre operativen Behandlungsmöglichkeiten. *Fortsch Kiefer Gesichtschir.* Bd 1, Thieme Verlag, Stuttgart (1955).
38. Watted, N., Bill, J.: Untergesichtsverlängerung bei der Klasse-II-Dysgnathien: Neues von der kieferorthopädisch-kieferchirurgischen Therapie. *Ästhetische Zahnmedizin* 4, 292-300 (1999).
39. Watted, N., Bill, J., Witt E., Reuther, J.: Lengthening of the lower face Angle class II patients with skeletally deep bite (short-face-syndrome) through combined orthodontic-surgical treatment. 75th Congress of the European Orthodontic Society Strasbourg, France (1999).

40. Watted, N., Teuscher, T., Wieber, M.: Die Analyse des vertikalen Gesichtaufbaus als Planungsorientierung in der orthognathen Chirurgie. Kieferorthop 16:29-44, (2002).
41. Watted, N., Witt, E., Bill, J.: Therapy Concept for the combined orthodontic-surgical Treatment of Angle Class II Deformities with Short-face Syndrome: New Aspects for surgical Lengthening of the Lower Face. Clin Orthod research3, 78-93 (2000).
42. Williamson, E.H., Caves, S.A., Edenfield, R.J., Morse, P.K.: Cephalometric analysis: comparisons between maximum intercuspatation and centric relation, Am J Orthod 74, 672-677 (1978).
43. Williamson, E.H., Evans, D. L., Barton, W.A., Williams B.H.: The effect of bite plane use on terminal hinge axis location. Angle Orthod 47, 25-33 (1977).
44. Westermark, A.H., Bystedt, H., von Konow, L., Sällström, K.O.: Nasolabial morphology after Le Fort I Osteotomies. Int J Oral Maxillofac Surg 20, 25-30, (1991).