

**Ausgabe:** Endodontie Journal 4/2017  
**Thema:** Die dreidimensionale endodontische Behandlungsplanung  
**Autor:** Dr. Jörg Tchorz

---

1. Dahlström L, Lindwall, Rystedt H, Reit C. 'Working in the dark': Swedish general dental practitioners on the complexity of root canal treatment. *Int Endod J.* 2017;50:636-645.
2. Sherwood IA. Pre-operative diagnostic radiograph interpretation by general dental practitioners for root canal treatment. *Dentomaxillofac Radiol.* 2012;41:43-54.
3. Tamse A, Kaffe I, Fishel D. Zygomatic arch interference with correct radiographic diagnosis in maxillary molar endodontics. *Oral Surg Oral Med Oral Pathol* 1980;50:563-566.
4. Uraba S, Ebihara A, Komatsu K, Ohbayashi N, Okiji T. Ability of Cone-beam Computed Tomography to Detect Periapical Lesions That Were Not Detected by Periapical Radiography: A Retrospective Assessment According to Tooth Group. *J Endod* 2016;42:1186-90.
5. Sousa TO, Haiter-Neto F, Nascimento EHL, Peroni LV, Freitas DQ, Hassan B. Diagnostic Accuracy of Periapical Radiography and Cone-beam Computed Tomography in Identifying Root Canal Configuration of Human Premolars. *J Endod.* 2017;43:1176-1179.
6. Mota de Almeida FJ, Knutsson K, Flygare L. The effect of cone beam CT (CBCT) on therapeutic decision-making in endodontics. *Dentomaxillofac Radiol* 2014;43:20130137.
7. Rodríguez G, Abella F, Durán-Sindreu F, Patel S, Roig M. Influence of Cone-beam Computed Tomography in Clinical Decision Making among Specialists. *J Endod* 2017;43:194-199.
8. Tchorz JP. 3D Endo: three-dimensional endodontic treatment planning. *Int J Comput Dent* 2017;20:87-92.