

**Ausgabe:** KN 12-2017, S. 1ff.  
**Thema:** KFO und Chirurgie  
**Autoren:** Prof. Dr. Nezar Watted, Prof. Dr. Dr. Peter Proff,  
Dr. Muhamad Abu-Hussein, Ali Watted

---

## Literatur

1.  
Helm, S., Siersbaek-Nielsen, S., Skieller, V., Björk, A.: Skelatal maturation of the hand in relation to maximum puberal growth in body height. Danish Dental Journal 75, 1223-1234 (1971).
2.  
Bass, N.M.: Dento-facial orthopaedics in der correction of the skeletal II malocclusion. Br J Orthod 9, 3-8 (1982).
3.  
Dryland-Vig, K.W.L., Ellis III, E.: Diagnosis and treatment planning for the surgical-orthodontic Patient. Cli Plast Surg 16, 645-658 (1989).
4.  
McNamara, J.A., McDougall, Jr.P.D., Dierks, J.M.: Arch with development in Class II patients treated with extraoral force and functional jaw orthodontics. Am J Orthodont 52, 353-359 (1966).
5.  
Petrovic, A.G., Stutzmann, J.: Reaktionsfähigkeit des tierischen und menschlichen Kondylenknorpels auf Zell- und Molekularebene im Lichte einer kybernetischen Auffassung des faszialen Wachstums. Fortschr Kieferorthop 49, 405-425 (1988).
6.  
Watted, N., Witt, E.: NMR study of TNJ changes following functional orthopaedic treatment using the "Würzburg approach", European Orthodontic Society (EOS) 74 th Congress (1998).
7.  
Watted, N.: Behandlung von Klasse II-Dysgnathien – Funktionskieferorthopädische Therapie unter besonderer Berücksichtigung der dentofazialen Ästhetik, Kieferorthop 13, 193-208 (1999).
8.  
Watted, N., Bill, J., Witt, E.: Therapy Concept for the Combined Orthodontic-Surgical Treatment of Angle Class II Deformities with Short Face Syndrome New Aspects for Surgical Lengthening of the Lower Face. Clin. Orthod. Res. 3, 78-93 (2000).
9.  
Watted, N., Bill, J., Witt, E., Reuther, J.: Lengthening of the lower face Angle class II patients with skeletally deep bite (short-face-syndrome) through combined orthodontic-surgical treatment. 75th Congress of the European Orthodontic Society Strasbuorg, France (1999).

10.

Watted, N., Teuscher, T., Wieber, M.: Vertikaler Gesichtsaufbau und Planung kieferorthopädisch-kieferchirurgischer Kombinationsbehandlungen unter besonderer Berücksichtigung der dentofazialen Ästhetik. *Kieferorthop* 16, 29-44 (2000).

11.

Witt, E.: Behandlungskonzepte. In Miethke, R.R., D. Drescher (Hrsg.): *Kleines Lehrbuch der Angel-Klasse II,1* unter besonderer Berücksichtigung der Behandlung. Quintessenz, Berlin (1996).

12.

Flanary, C.M., Barnwell, G.M., Alexander, J.M.: Patient perceptions of orthognathic surgery. *Am J Orthod* 88, 137-145 (1985).

13.

Jacobson, A.: The influence of children's dentofacial appearance on their social attractiveness as judged by peers and lay adults. *Am J Orthod* 79, 399-415 (1981).

14.

Kiyak, H.A., Hohl, T., West, R.A.: Psychologie changes in orthognathic surgery patients: a 24-month follow-up. *J Oral Maxillofac Surg* 42, 506-512 (1984).

15.

Albino, J.E., Tedesco, L.: Esthetic need for orthodontic treatment. In Melsen B, editor: *Current controversies in orthodontics*. Chicago, , Quintessence Publishing, pp. 11-24 (1994).

16.

Berscheid, E., Gangestad, S: The social psychological implications of facial physical attractiveness. *Clin Plast Surg* 9, 289-296 (1982).

17.

Scott, O., Kijak, H.A.: Treatment expectation versus outcomes among orthodontic surgery patients. *Int J Adult Orthod Orthognath Surg* 6, 247-255 (1991).

18.

Canut, J.: Eine Analyse der dentofazialen Ästhetik. *Inf Orthod Kieferorthop* 28, 83-105 (1996).

19.

Farkas, L.G., Kolar, J.C.: Anthropometry and art in the aesthetics of women's face. *Clin Plast Surg* 14, 599-615 (1987).

20.

Legan, H.L., Burstone, G.J.: Soft tissue cephalometric analysis for orthognathic surgery. *J Oral Surg* 38, 744-51 (1980).

21.

Witt, E.: Möglichkeiten und Grenzen der kieferorthopädischen Behandlung Erwachsener. *Fortschr Kieferorthop* 52, 1-7 (1991).

22.

Westermarck AH, Bystedt H, von Konow L, Sällström KO. Nasolabial morphology after Le Fort I Osteotomies. *Int J Oral Maxillofac Surg* 1991; 20: 25-30.

23.

Wolford L.M., Epker B.N.: The combined anterior and posterior maxillary osteotomy a new technique. *J. Oral Surg.* 14: 123-127, 1975.

24.

Williamson EH, Evans DL, Barton WA, Williams BH. The effect of bite plane use on terminal hinge axis location, *Angle Orthod.* 1977; 47: 25-33.

25.

Williamson EH, Caves SA, Edenfield RJ, Morse PK. Cephalometric analysis: comparisons between maximum intercuspation and centric relation, *Am J Orthod* 1978; 74: 672-677.

26.

Schwarz, A.M.: *Die Röntgendiagnostik.* Urban & Schwarzenberg, Wien (1958)