

Ausgabe: Cosmetic Dentistry 1/2018

Thema: Amelogenesis imperfecta. Behandlung einer Funktionsstörung
mittels Bisslagenveränderung

Autoren: Dr. Sven Egger; ZTM Christian Berg

Literatur

- 1 Rufenacht Claudia, R.: Fundamentals of Esthetics. Chicago, Quintessenz, 1990; 67-134.
- 2 Magne Pascal, Belser Urs: Adhäsiv befestigte Keramikrestaurationen; Berlin, Quintessenz, 2004
- 3 Fradeani Mauro: Evaluation of dentolabial parameters as part of a comprehensive esthetic analysis; Eur J Esthet Dent. 2006; 1; 62-69.
- 4 D'Amico Angelo: The canine teeth-normal functional relation of the natural teeth of man; J South Calif Dent Assoc. 1958; 26: 6-23, 49-60, 127-142, 175-182, 194-208, 239-241.
- 5 Magne Pascal, Douglas William H.: Additive contour of porcelain veneers: A key element in enamel preservation, adhesion and esthetic for the aging dentition; J Adhesive Dent 1999; 1; 81-91.
- 6 Vailati Francesca, Belser Urs Christoph: Palatal and facial veneers to treat severe dental erosion: a case report following the three-step technique and the sandwich approach; Eur J Esthet Dent 2011; Autumn, 6(3); 268-78.
- 7 Witkowski Siegbert, Schicha Kerstin: Prep Veneers & Non-Prep Veneers; Berlin; Quintessenz, 2010.
- 8 Peumans M, Van Meerbeek B, Lambrechts P, Vanherle G, Quirynen M: The influence of direct composite additions for the correction of tooth form and/or position on periodontal health: A retrospective stud.; J Periodontol 1998; 69; 422-427.