

## Literaturliste

### Einsatz von Laser bei Sofortimplantation

*Dr. Michael Schäfer/Düsseldorf*

*Laser Journal 3/2011*

Araujo MG, Lindhe J: Dimensional ridge alterations following tooth extraction. An experimental study in the dog. J Clin Periodontol 32: 212-218 (2005)

Atwood, D. A.: Reduction of residual ridges: A major oral disease entity. J Prosthe Dent 26, 266-279 (1971)

Boticelli D, Berglundh T, Lindhe J: Hard-tissue alterations following immediate implant placement in extraction sites. J Clin Periodontol 8: 820-828 (2004)

Chen ST, Wilson Jr. TG, Hämmerle CH: Immediate or early placement of implants following tooth extraction: review of biological basis, clinical procedures and outcomes. Int J Oral Maxillofac Implants 19 (Suppl):12-25 (2004)

Chen S T, Darby I B, Reynolds E C: Prospective clinical study of non-submerged immediate implants: clinical outcomes and esthetic results. Clin Oral Implants Res. 18: 552–562 (2007)

Geistlich Infobrochure: Therapiekonzepte für die Extraktionsalveole. 2008

Grunder U, Gracis S, Capelli M.: Influence of the three dimensional bone-to-implant relationship on esthetics  
Int J Periodont Rest Dent 2005; 25:113-119

Hämmerle Ch, Chen S, Wilson Th: Consensus statement and recommended clinical procedure regarding the placement of implants in extraction sockets. Int J Oral Maxillofac Implants 19: 26-28 (2008)

Kan JY, Rungcharassaeng K, Umezue K, Kois JC: Dimensions of peri-implant mucosa: an evaluation of maxillary anterior single implants in humans. J Periodontol 74 (4): 557-562 (2003)

Schropp L, Wenzel A, Kostopoulos L, Karring T: Bone healing and soft tissue contour changes following single-tooth extraction. A clinical and radiographic 12-month prospective study. Int J Periodontics Restorative Dent 23:312-323 (2003)

Tarnow DP: The Effect of Inter-Implant Distance on the Height of the presence or absence of the interproximal dental papilla. J Periodontol 1992; 63:995-996.