

Ausgabe: Cosmetic Dentistry 3/2018

Thema: Einzelzahnveneer in der Oberkieferfront – eine Herausforderung

Autoren: Dr. Andreas Koch

Literatur

1. Beier US, Kapferer I, Burtscher D, Dumfahrt H. Clinical performance of porcelain laminate veneers for up to 20 years. *Int J Prosthodont.* 2012;25:79–85.
2. Layton DM, Clarke M. A systematic review and meta-analysis of the survival of non-feldspathic porcelain veneers over 5 and 10 years. *Int J Prosthodont.* 2013 Mar-Apr;26(2):111-24. doi: 10.11607/ijp.3202.
3. Morimoto S, Albanesi RB, Sesma N, Agra CM, Braga MM. Main Clinical Outcomes of Feldspathic Porcelain and Glass-Ceramic Laminate Veneers: A Systematic Review and Meta-Analysis of Survival and Complication Rates. *Int J Prosthodont.* 2016 Jan-Feb;29(1):38-49. doi: 10.11607/ijp.4315.
4. Granell-Ruiz M, Fons-Font A, Labaig-Rueda C, Martinez-Gonzalez A, Roman-Rodriguez JL, Sola-Ruiz MF. A clinical longitudinal study 323 porcelain laminate veneers. Period of study from 3 to 11 years. *Med Oral Patol Oral Cir Bucal* 2010;15:e531-e537.
5. Piemjai M, Arksornnukit M. Compressive fracture resistance of porcelain laminates bonded to enamel or dentin with four adhesive systems. *J Prosthodont* 2007; 16:457-464.
6. Peumans M, Van Meerbeek B, Lambrechts P, Vanherle G. Porcelain veneers: A review of the literature. *J Dent* 2000; 28:163-177
7. Igiel C, Weyhrauch M, Mayer B, Scheller H, Lehmann KM. Effects of ceramic layer thickness, cement color, and abutment tooth color on color reproduction of feldspathic veneers. *Int J Esthet Dent.* 2018;13(1):110-119.