

Ausgabe: Endodontie Journal 3/2018

Thema: Apikale Läsion – behandeln oder nicht?

Autor: Dr. Christophe Verbanck

Literatur

1 Kim D, Ku H, Nam T, Yoon TC, Lee CY, Kim E. Influence of size and volume of periapical lesions on the outcome of endodontic microsurgery: 3 dimensional analysis using cone beam computed tomography. J endod 2016, Aug 42 (8) 1196- 1201.

2 Kruse C, Spin-Neto R, Wenzel A, Kirkevang LL. Cone beam computed tomography and periapical lesions: a systematic review analysing studies on diagnostic efficacy by a hierarchical model. Int endod J, 2015, Sep48(9), 815-28.

3 Morris MF, Kirkpatrick TC, Rutledge RE, Schindler WG. Comparison of non surgical root canal treatment and single tooth implants. J endod 2009 oct 35 (10), 1325-30.

4 Neukermans M., Vanobbergen J., De Bruyne M., Meire M., De Moor R.J.G. Endodontic performance by Flemish dentists: have they evolved? International Endodontic Journal, 2014.

5 Ng YL, Mann V, Gulabivala K. A prospective study of the factors affecting outcomes of nonsurgical root canal treatment: part 1: periapical health. Int endod J. 2011, Jul, 44(7), 583-609.

6 P.N.R Naïr. Review New perspectives on radicular cysts: do they heal? International endodontic journal, 1998, 31, 155-160.

7 Sundqvist G. Bacteriological studies of necrotic dental pulps. Umeå Univ Odontological Dissertations 1976.

8 Wu MK, Shemesh H, Wesselink PR. Limitations of previously published systematic reviews evaluating the outcome of endodontic treatment. Int endod J. 2009, Aug 42(8), 656-66.

9 Yu VS, Khin LW, Hsu CS, Yee R, Messer HH. Risk score algorithm for treatment of persistent apical periodontitis. J dent Res, 2014, nov, 93(11); 1076-82.