

Ausgabe: KN 9-2018, S. 1ff.

Thema: Gesicht-Okklusion-Atmung

Autoren: Priv.-Doz. Dr. Dr. Robin Seeberger

Literatur

- 1: Alves PV, Zhao L, O'Gara M et al (2008) Three-dimensional cephalometric study of upper airway space in skeletal Class II and III healthy patients. *J Craniofac Surg* 19:1497–1507
- 2: Park JW, Kim NK, Kim JW et al (2010) Volumetric, planar, and linear analyses of pharyngeal airway change on computed tomography and cephalometry after mandibular setback surgery. *Am J Orthod Dentofacial Orthop* 138:292–299
- 3: Schutz TC, Dominguez GC, Hallinan MP et al (2011) Class II correction improves nocturnal breathing in adolescents. *Angle Orthod* 81:222–228
- 4: Stellzig-Eisenhauer A, Meyer-Marcotty P (2010) Interaction between otorhinolaryngology and orthodontics: correlation between the nasopharyngeal airway and the craniofacial complex. *GMS Curr Top Otorhinolaryngol Head Neck Surg* 9:Doc04
- 5: Kochel J, Meyer-Marcotty P, Sickel F, Lindorf H, Stellzig-Eisenhauer A. (2013) Short-term pharyngeal airway changes after mandibular advancement surgery in adult Class II-Patients—a three-dimensional retrospective study. *J Orofac Orthop.* 2013 Mar;74(2):137-52.
- 6: Haas, A.J., Palatal expansion: just the beginning of dentofacial orthopedics. *Am J Orthod*, 1970. 57(3): p. 219-55.
- 7: Seeberger R, Gander E, Hoffmann J, Engel M.: Surgical management of cross-bites in orthognathic surgery: Surgically assisted rapid maxillary expansion (SARME) versus two-piece maxilla. *J Craniomaxillofac Surg.* 2015 Sep;43(7):1109-12.
- 8: Seeberger R, Kater W, Schulte-Geers M, Thiele OC, Davids R, Hofele CH, et al. [Surgically assisted rapid maxillary expansion. Effects on the nasal airways and nasal septum]. *Hno.* 2010;58(8):806-11
- 9: Seeberger R, Kater W, Schulte-Geers M, Davids R, Freier K, Thiele O: Changes after surgically-assisted maxillary expansion (SARME) to the dentoalveolar, palatal and nasal structures by using tooth-borne distraction devices. *Br J Oral Maxillofac Surg.* 2011 Jul;49(5):381-5.
- 10: Seeberger R, Abe-Nickler D, Hoffmann J, Kunzmann K, Zingler S.: One-stage tooth-borne distraction versus two stage bone-borne distraction in surgically assisted maxillary expansion (SARME). *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2015 Dec;120(6):693-8.
- 11: Tausche E, Deeb W, Hansen L, Hietschold V, Harzer W, Schneider M (2009): CT analysis of nasal volume changes after surgically-assisted rapid maxillary expansion. *J Orofac Orthop;*70(4):306-17
- 12: Timms DJ (1984): The reduction of nasal airway resistance by rapid maxillary expansion and its effect on respiratory disease. *J Laryngol Otol.*; 98(4): 357-62
- 13: Bell WH, Epker BN. Surgical-orthodontic expansion of the maxilla. *American journal of orthodontics.* 1976;70(5):517-28.

- 14: Obwegeser, H., The Indications for Surgical Correction of Mandibular Deformity by the Sagittal Splitting Technique. *Br J Oral Surg*, 1964. 1: p. 157-71.
- 15: Hunsuck EE. A modified intraoral sagittal splitting technic for correction of mandibular prognathism. *J Oral Surg*. 1968;26:250–253.
- 16: Epker BN. Modifications in the sagittal osteotomy of the mandible. *J Oral Surg*. 1977;35:157–159.
- 17: Seeberger R, Asi Y, Thiele OC, Hoffmann J, Stucke K, Engel M.: Neurosensory alterations and function of the temporomandibular joint after high oblique sagittal split osteotomy: an alternative technique in orthognathic surgery. *Br J Oral Maxillofac Surg*. 2013 Sep;51(6):536-40.
- 18: Kuehle R, Berger M, Saure D, Hoffmann J, Seeberger R.: High oblique sagittal split osteotomy of the mandible: assessment of the positions of the mandibular condyles after orthognathic surgery based on cone-beam tomography. *Br J Oral Maxillofac Surg*. 2016 Jul;54(6):638-42. doi: 10.1016/j.bjoms.2016.03.017. Epub 2016 Apr 3.
- 19: Zingler S, Hakim E, Finke D, Brunner M, Saure D, Hoffmann J, Lux CJ, Erber R, Seeberger R.: Surgery-first approach in orthognathic surgery: Psychological and biological aspects - A prospective cohort study. *J Craniomaxillofac Surg*. 2017 Jun 6.
- 20: Proffit WR, Miguel JA: The duration and sequencing of surgical-orthodontic treatment. *Int J Adult Orthodon Orthognath Surg* 10:35, 1995.
- 21: Alves PV, Zhao L, O'Gara M et al (2008) Three-dimensional cephalometric study of upper airway space in skeletal Class II and III healthy patients. *J Craniofac Surg* 19:1497–1507
- 22: Schutz TC, Dominguez GC, Hallinan MP et al (2011) Class II correction improves nocturnal breathing in adolescents. *Angle Orthod* 81:222–228
- 23: Caples SM, Rowley JA, Prinsell JR, Pallanch JF, Elamin MB, Katz SG, Harwick JD. (2010) Surgical modifications of the upper airway for obstructive sleep apnea in adults: a systematic review and meta-analysis. *Sleep*. 2010 Oct;33(10):1396-407.
- 24: Mehra P, Downie M, Pita MC et al (2001) Pharyngeal airway space changes after counterclockwise rotation of the maxillomandibular complex. *Am J Orthod Dentofacial Orthop* 120:154–159