

Ausgabe: Prophylaxe Journal 5/2018

Thema: Mundtrockenheit bedeutet erhöhtes Risiko für die Zähne

Autor: Dr. Gabriele David

Literatur

1. Plemons JM, Al-Hashimi I, Marek CL, American Dental Association Council on Scientific Affairs: Managing xerostomia and salivary gland hypofunction. JADA 2014; 145(8): 867-873.
2. Edgar WM, O`Mullane DM (Hrsg): Saliva and oral health. BDJ 1996; Second Edition.
3. Turner MD, Ship JA: Dry mouth and its effects on the oral health of elderly people. JADA 2007; 15S-20S.
4. De Paola PF, Soparkar P, Foley S, Bookstein F, Bakhos Y: Effect of high-concentration ammonium and sodium fluoride rinses on dental caries in schoolchildren. Community Dent Oral Epidemiol 1977; 5: 7-14.
5. Fischer K: Wissenschaftliche Dokumentation Fluor Protector S. Ivoclar Vivadent 2013.
6. Fischer K: Wissenschaftliche Dokumentation Fluor Protector Gel, Ivoclar Vivadent 2010.
7. Al-Mashhadani A, Plygkos I, Bozec L, Rodriguez JM: Three-dimensional in vitro measurements of tooth wear using fluoridated dentifrices. Aust Dent J 2016; 61(3): 304-309.