

**Ausgabe:** KN 11-2018 (Teil 1) / KN 12-2018 (Teil 2), S. 10ff.

**Thema:** Hypodontie-Dienstleistungen des National Health Service  
in Großbritannien

**Autoren:** Dr. Dirk Bister

---

## **Literatur**

Abu-Awwad, Hemmings K, Mannaas, Gill D, GulamaliA, Petrie A.: Treatment Outcomes and Assessment of Oral Health Related Quality of Life in Treated Hypodontia Patients. EurJ ProsthodontRestorDent. 2017 Mar; 25(1):49-56.

Bjerkin K., Bennet J.: The long-term survival of lower second primary molars in subjects with agenesis of the premolars. EJO 2000 (22): 245-255.

Dunne S. M., Millar B. J.: A longitudinal study of the clinical performance of resin bonded bridges and splints. BDJ: 1993; 174: 405-411.

Garnett et al.: Survival of resin-bonded bridgework provided for post orthodontic hypodontia patients with missing maxillary lateral incisors, BDJ 201 (8) 527 –534; 2006.

Luck O, Harzer W, Boening K: Orthodontic treatment planning for single tooth implants in patients with congenitally missing lateral incisors. EOS Congress, June 2003, Prague.

Mall, C; Schneider, E; Jonas, I E: Orthodontic versus implant prosthetic space closure in subjects with missing upper incisors. EOS Congress, June 2003, Prague

Millar B.J.; Taylor N.G.: Lateral thinking: the management of missing upper lateral incisors. BDJ 1995, August: 99 -106.

Nordquist, McNeill: Orthodontic vs. Restorative Treatment of the Congenitally Absent Lateral Incisor -Long term Periodontal and Occlusal Evaluation. J. Periodontology 1975; 46 (3), 139 -143.

Spear F.M., Mathews D.M., Kokich V.G.: Interdisciplinary Management of Single -Tooth implants. Seminars in Orthodontics, 1997 Vol 3 (1); 45-72.

Thoradson A., Zachrisson B.U., Mjor I.A.: Remodeling of canines to the shape of lateral incisors by grinding: a long -term clinical and radiographic evaluation. AJODO 1991; 100: 123 -132.