

LITERATUR

Ausgabe: Implantologie Journal 5/19

Thema: Kieferkammaugmentationen – drei Wege, ein Ziel
Neue minimalinvasive Goldstandards

Autor: OA Dr. Dr. Angelo Troedhan, v. Prof., MD, DMD, PhD

[1] Reich, K. M., Huber, C. D., Lippnig, W. R., Ulm, C., Watzek, G., & Tangl, S. (2011). Atrophy of the residual alveolar ridge following tooth loss in an historical population. *Oral Diseases*, 17(1), 33-44.

[2] Bras J (1990). Mandibular Atrophy and Metabolic Bone Loss. *International Dental Journal* 40: 298-302.

[3] Cawood, J. I., & Howell, R. A. (1988). A classification of the edentulous jaws. *International journal of oral and maxillofacial surgery*, 17(4), 232-236.

[4] Cordaro, L., Amadè, D. S., & Cordaro, M. (2002). Clinical results of alveolar ridge augmentation with mandibular block bone grafts in partially edentulous patients prior to implant placement. *Clinical oral implants research*, 13(1), 103-111.

[5] Buser, D., Dula, K., Hirt, H. P., & Schenk, R. K. (1996). Lateral ridge augmentation using autografts and barrier membranes: a clinical study with 40 partially edentulous patients. *Journal of oral and maxillofacial surgery*, 54(4), 420-432.

[6] Sanz-Sanchez, I., Ortiz-Vigon, A., Sanz-Martin, I., Figuero, E., & Sanz, M. (2015). Effectiveness of lateral bone augmentation on the alveolar crest dimension: a systematic review and meta-analysis. *Journal of dental research*, 94(9_suppl), 128S-142S.

[7] Silva, F. M. S., Cortez, A. L. V., Moreira, R. W. F., & Mazzone, R. (2006). Complications of intraoral donor site for bone grafting prior to implant placement. *Implant dentistry*, 15(4), 420-426.

[8] Troedhan, A., Kurrek, A., & Wainwright, M. (2012). Vertical alveolar crest split and widening—an experimental study on cow ribs, ultrasonic tool development and test on human cadaver heads. *Surgical Techniques Development*, 2(1), 10.

- [9] Troedhan, A., Kurrek, A., Wainwright, M., & Schlichting, I. (2015). Flapless Piezotome-Enhanced Vertical Alveolar Crest-Split and Horizontal Distraction of Alveolar Crests (FPeCSWT) of Less than 2 mm Width: Results of a Prospective Comparative 3-Year Clinical Multicenter-Study with 239 Patients, 261 Crest-Split Sites and 488 Inserted Dental Implants. *Open Journal of Stomatology*, 5(07), 159.
- [10] Troedhan, A., Tarek Mahmoud, Z., & Wainwright, M. (2018). Piezotome-crest-split versus buccal autologous onlay grafts-results of a randomized clinical trial. *Clinical Oral Implants Research*, 29, 92-92.
- [11] Rothstein, S. S., Paris, D. A., Zacek, M. P. (1984). Use of hydroxylapatite for the augmentation of deficient alveolar ridges. *Journal of oral and maxillofacial surgery*, 42(4), 224-230.
- [12] Propper, R. H. (1985). A technique for controlled placement of hydroxylapatite over atrophic mandibular ridges. *Journal of oral and maxillofacial surgery*, 43(6), 469-470.
- [13] Härle, F., Kreuzsch, T. (1991). Augmentation of the alveolar ridges with hydroxylapatite in a Vicryl tube. *International journal of oral and maxillofacial surgery*, 20(3), 144-148.
- [14] Troedhan, A., Wainwright, M., Kurrek, A., Schlichting, I. (2015). Biomechanical stability of dental implants in augmented maxillary sites: results of a randomized clinical study with four different biomaterials and PRF and a biological view on guided bone regeneration. *BioMed research international*, 2015.
- [15] Spin-Neto, R., Stavropoulos, A., Coletti, F. L., Pereira, L. A., Marcantonio Jr, E., & Wenzel, A. (2015). Remodeling of cortical and corticocancellous fresh-frozen allogeneic block bone grafts—a radiographic and histomorphometric comparison to autologous bone grafts. *Clinical oral implants research*, 26(7), 747-752.
- [16] Schmidlin, P. R., Nicholls, F., Kruse, A., Zwahlen, R. A., & Weber, F. E. (2013). Evaluation of moldable, in situ hardening calcium phosphate bone graft substitutes. *Clinical oral implants research*, 24(2), 149-157.
- [17] Kalfas, I. H. (2001). Principles of bone healing. *Neurosurgical focus*, 10(4), 1-4.