

Ausgabe: ZWP Zahnarzt Wirtschaft Praxis 7+8/2019

Thema: Die präendodontische Phase

Autor: Dr. Preet Singh

Literatur

1 Kakehashi, S., H. R. Stanley and R. J. Fitzgerald (1965). "The Effects of Surgical Exposures of Dental Pulps in Germ-Free and Conventional Laboratory Rats." *Oral Surg Oral Med Oral Pathol* 20: 340-349.

2 Jordan, R. E., M. Suzuki and D. H. Skinner (1978). "Indirect pulp-capping of carious teeth with periapical lesions." *J Am Dent Assoc* 97(1): 37-43.

3 Sato, T., E. Hoshino, H. Uematsu and T. Noda (1993). "In vitro antimicrobial susceptibility to combinations of drugs on bacteria from carious and endodontic lesions of human deciduous teeth." *Oral Microbiol Immunol* 8(3): 172-176.

4 Marsh P, M. M. (2009). *Oral Microbiology*. Oxford, Elsevier Health Sciences.

5 Mejare, B. (1975). "Streptococcus salivarius, Streptococcus sanguis and Streptococcus "mitis" as indicators of contamination of root canal cultures." *Odontol Revy* 26(2): 125-134.

6 https://www.dgzmk.de/uploads/tx_szdgzmkdocuments/Good_clinical_practice:_Die_Wurzel_kanalbehandlung.pdf

7 Ray, H. A. and M. Trope (1995). "Periapical status of endodontically treated teeth in relation to the technical quality of the root filling and the coronal restoration." *Int Endod J* 28(1): 12-18.