

## LITERATUR

**Ausgabe:** Oralchirurgie Journal 3/19

**Thema:** Konzept für kieferorthopädisch-kieferchirurgische Therapie –  
Behandlung skelettaler Dysgnathien Klasse III

**Autoren:** Prof. Dr. med. dent. Nezar Watted, Abu-Hussein Muhamad, Obeida  
Awadi, Omri Emodi, Shelo Dekel, Prof. Dr. med. Dr. med. dent. Peter  
Proff, Ali Watted

---

- 1) Albino, J.E., Tedesco, L.: Esthetic need for orthodontic treatment. In Melsen B, editor: Current controversies in orthodontics. Chicago, , Quintessence Publishing, pp. 11-24 (1994).
- 2) Al-Riyami S, Cunningham SJ, Moles DR. (2009) Orthognathic treatment and temporomandibular disorders: a systematic review. Part 2. Signs and symptoms and meta-analyses. Am J Orthod Dentofacial Orthop. 136(5):626.e1-16, discussion 626-7.
- 3) Bass, N.M.: Dento-facial orthopaedics in der correction of the skeletal II malocclusion. Br J Orthod 9, 3-8 (1982).
- 4) Bell, H.W., Jacobs, J.D.: Simultaneous repositioning of the maxilla, mandible, and chin. Treatment planning and analysis of the soft tissues. Am J Orthod 91, 28-50 (1986).
- 5) Bell, W.H., Proffit, H.R., White, R.P.: Surgical correction of facial deformities. Saunder Verlag, Baltimor (1980).
- 6) Bell, W.H., Schdeman, G.B.: Correction of vertical maxillary deficiency: Stability and soft tissue changes. J Oral surg 39, 666-670 (1981).
- 7) Berscheid, E., Gangestade, S: The social psychological implications of facial physical attractiveness. Clin Plast Surg 9, 289-296 (1982).
- 8) Buddgaard, M., Melsen, B., Terp, S.: Changes during and following total maxillary osteotomy (Le Fort I Procedure): a cephalometric study. Eur J Orthod 8, 21-29 (1986).
- 9) Canut ,J.: Eine Analyse der dentofazialen Ästhetik. Inf Orthod Kieferorthop 28, 83-105 (1996).

- 10) Centenero SAH, Hernández-Alfaro F (2012) 3D planning in orthognathic surgery: CAD/CAM surgical splints and prediction of the soft and hard tissues results-Our experience in 16 cases. J Craniomaxillofac Surg 40: 162-168.
- 11) Collins, P., Epker, B.N.: The alar basse Cinch: A technique for prevention of alar base flaring secondary to maxillary surgery. Oral Surg 53, 549-554 (1982).
- 12) Dal Pont G.: L`osteotomia retromolare per la correzione della progenia, Minerva chir. 18, 1138-1141 (1959).
- 13) Dal Pont G.: Die retromolare Osteotomie zur Korrektur der Progenie, der Retrogenie und des Mordex apertus. Öst Z Stoma 58, 8-10 (1961).
- 14) Dryland-Vig, K.W.L., Ellis III, E.: Diagnosis and treatment planning for the surgical-orthodontic Patient. Cli Plast Surg 16, 645-658 (1989).
- 15) Dann, J.J., Fonseca, R.J., Bell, W.H.: Soft tissue changes associated with total maxillary advancement: a preliminary study. J Oral Surg 34, 19-23 (1976).
- 16) De Assis, E.A., Starck, W.J., Epker, B.: Cephalometric analysis of profile nasal esthetics. Part III. Postoperativ changes after isolated superior repositioning. Int J Adult Orthod Orthognath 11, 279-288 (1996).
- 17) Ellis, E.: Condylar Positioning Devices for Orthodontic Surgery: Are They Necessary. J Oral Maxillofac Syrg 52, 536-552 (1994).
- 18) Ellis, E., Hinton R.J.: Histologic examination of the temporomandibular joint after mandibular advancement with and without rigid fixation: An experimental investigation in adult Maccaca mulatta. J Oral Maxillofac Syrg 49, 1316 (1991).
- 19) Epker, B.: Esthetic maxillofacial surgery. Lea und Febiger Verlag, Philadelphia (1994).
- 20) Epker, B. N., Fish, L. C.: Surgical-orthodontic correction of open-bite deformity. Am J Orthod 71, 278-299 (1977).
- 21) Epker, B.N., Fish, L.C.: Surgical superior repositioning of the maxilla: What to do with the mandibile?. Am j Orthod 78, 164-191 (1980).
- 22) Farkas, L.G., Kolar, J.C.: Anthropometry and art in the aesthetics of women´s face. Clin Plast Surg 14, 599-615 (1987).

- 23) Flanary, C.M., Barnwell, G.M., Alexander, J.M.: Patient perceptions of orthognathic surgery. *Am J Orthod* 88, 137-145 (1985).
- 24) Freihofer, H.P.M.: Changes in nasal Profile after maxillary advancement in cleft and non-cleft patients. *J Maxillofac Surg* 5, 20-27 (1977).
- 25) Helm, S., Siersbaek-Nielsen, S., Skieller, V., Björk, A.: Skelatal maturation of the hand in relation to maximum puberal growth in body height. *Danish Dental Jornal* 75, 1223-1234 (1971).
- 26) Hogeman, K.E., Sarnäs, K.V., Willmar, K.: One-stage surgical and dental-orthopedic correction of bimaxillary and craniofacial dysostosis. In: *Fortschr Kiefer Gesichtschr, Bd 18: 39-49*, Thieme Verlag, Stuttgart (1974)
- 27) Hui, E., Hägg, E.U.O. Tidman, H: Soft tissue changes following maxillary osteotomies in cleft lip and palate and non-cleft Patients. *J Craniomaxillofac Surg* 22, 182-186 (1994).
- 28) Jensen, A.C., Sinclair, P.M., Wolford, L.M.: Soft tissue changes associated with double jaw surgery. *Am J Orthod Dentofac Orthop* 101, 266-275 (1992).
- 29) Jacobson, A.: The influence of children´s dentofacial appearance on their social attractiveness as judged by peers and lay adults. *Am J Orthod* 79, 399-415 (1981).
- 30) Joss CU, Vassalli IM. Stability after bilateral sagittal split osteotomy dvancement surgery with rigid internal fixation: a systematic review. *J Oral Maxillofac Surg.* 2009;67:301-13
- 31) Kiyak, H.A., Hohl ,T., West, R.A.: Psychologie changes in orthognathic surgery patients: a 24-month follow-up. *J Oral Maxillofac Surg* 42, 506-512 (1984).
- 32) Lazaridou-Terzoudi T, Kiyak HA, Moore R, Athanasiou AE, Melsen B (2003) Long-term assessment of psychologic outcomes of orthognathic surgery. *J Oral Maxillofac Surg* 61:545–552
- 33) Legan, H.L., Burstone, G.J.: Soft tissue cephalometric analysis for orthognathic surgery. *J Oral Surg* 38, 744-51 (1980).
- 34) Lee, D.Y., Bailey, L.J., Proffit, W.R.: Soft tissue changes after superior repositioning of the Maxilla with Le Fort I osteotomy: 5-years follow-up. *Int J Adult Orthod Orthognath* 11, 301-311 (1996).

- 35) McNamara, J.A., McDougall, Jr.P.D., Dierks, J.M.: Arch with development in Class II patients treated with extraoral force and functional jaw orthodontics. *Am J Orthodont* 52, 353-359 (1966).
- 36) Mensink G, Verweij JP, Frank MD, Eelco BJ, Richard van Merkesteyn JP. Bad split during bilateral sagittal split osteotomy of the mandible with separators: a retrospective study of 427 patients. *Br J Oral Maxillofac Surg*. 2013;51:525–529.
- 37) Mensink G, Zweers A, Wolterbeek R, Dicker GG, Groot RH, van Merkesteyn RJ. Neurosensory disturbances one year after bilateral sagittal split osteotomy of the mandibula performed with separators: A multi-centre prospective study. *J Craniomaxillofac Surg*. 2012;40:763–767.
- 38) Obwegeser, H.: The surgical correction of mandibular prognathism and retrognathia with consideration of genioplasty. *J Oral Surg* 10, 677-689 (1957).
- 39) Obwegeser, H.: The indication for surgical correction of mandibular deformity by sagittal splitting technique. *Br J Surg* 1, 157-168 (1963).
- 40) Obwegeser, H., Trauner, R.: Zur Operationstechnik bei der Progenie und anderen Unterkieferanomalien. *Dtsch Zahn Mund Kieferheilk* 23, 1-26 (1955).
- 41) Petrovic, A.G., Stutzmann, J.: Reaktionsfähigkeit des tierischen und menschlichen Kondylenknorpels auf Zell- und Molekularebene im Lichte einer kybernetischen Auffassung des faszialen Wachstums. *Fortschr Kieferorthop* 49, 405-425 (1988).
- 42) Proothi M, Drew SJ, Sachs SA. Motivating factors for patients undergoing orthognathic surgery evaluation. *J Oral Maxillofac Surg*. 2010;68:1555–1559
- 43) Radney, L.J., Jacobs, J.D.: Soft tissue changes associated with surgical total maxillary intrusion. *Am J Orthodont* 80, 191-212 (1981)
- 44) Rosen, H.M.: Lip-nasal esthetics follwing Le Fort I Osteotomy. *Plast Reconstr Surg* 81, 171-182 (1988)
- 45) Schwarz, A.M.: Die Röntgendiagnostik. Urban & Schwarzenberg, Wien (1958).
- 46) Scott, O., Kijak, H.A.: Treatment expectation versus outcomes among orthodontic surgery patients. *Int J Adult Orthod Orthognath Surg* 6, 247-255 (1991).

- 47) Solano-Hernandez B, Antonarakis GS, Scolozzi P, Kiliaridis S. (2013) Combined orthodontic and orthognathic surgical treatment for the correction of skeletal anterior open-bite malocclusion: a systematic review on vertical stability. J Oral Maxillofac Surg 71(1):98-109.
- 48) Watted, N., Witt, E.: NMR study of TNJ changes following functional orthopaedic treatment using the "Würzburg approach", European Orthodontic Society (EOS) 74 th Congress (1998).
- 49) Watted, N.: Behandlung von Klasse II-Dysgnathien-Funktionskieferorthopädisch Therapie unter besonderer Berücksichtigung der dentofazialen Ästhetik, Kieferorthop 13, 193-208 (1999).
- 50) Watted, N., Bill, J., Witt, E.:Therapy Concept for the Combined Orthodontic-Surgical Treatment of Angle Class II Deformities with Short Face Syndrome New Aspects for Surgical Lengthening of the Lower Face. Clinc. Orthod. Res. 3, 78-93 (2000).
- 51) Watted, N., Bill, J., Witt, E., Reuther, J.: Lengthening of the lower face Angle class II patients with skelettaly deep bite (short-face-syndrome) through combined orthodontic-surgical treatment. 75th Congress of the European Orthodontic Society Strasbuorg, France(1999).
- 52) Watted, N., Teuscher, T., Wieber, M.:Vertikaler Gesichtsaufbau und Planung kieferorthopädisch-kieferchirurgischer Kombinationsbehandlungen unter besonderer Berücksichtigung der dentofazialen Ästhetik. Kieferorthop 16, 29-44 (2000).
- 53) Watted, N., Abu-Hussein, M., Bill, J., Proff, P Harmonisation of the Dento-Facial Complex A Result Of Combination A Orthodontic and Orthognathic Surgical Therapy. IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)14, 2: 128-139., (2015)
- 54) Watted, N., Abu-Hussein, M. Watted, A., Bill, J., Proff, P., Schlomi, B. Änderung der skelettalen Strukturen zur Harmonisierung der Weichteilstrukturen im Gesicht, Oralchirurgie Journal 12: 41-48 (2015)
- 55) Watted, N., Abu-Hussein, M, Hussein, E , Abu Mowais, M, P. Proff, Watted, A Ein integriertes Therapiekonzept zur Korrektur der Klasse-II-Dysgnathien durch kombinierte kieferorthopädisch-kieferchirurgische Behandlung- Part II, ZMK, 33, 9: 1-12, (2017)

- 56) Westermarck, A.H., Bystedt, H., von Konow, L., Sällström, K.O.: Nasolabial morphology after Le Fort I Osteotomies. *Int J Oral Maxillofac Surg* 20, 25-30, (1991).
- 57) Williamsone, E.H., Caves, S.A., Edenfield, R.J., Morse, P.K.: Cephalometric analysis: comparisons between maximum intercuspation and centric relation, *Am J Orthod* 74, 672-677 (1978).
- 58) Williamsone, E.H., Evans, D. L., Barton, W.A., Williams B.H.: The effect of bite plane use on terminal hinge axis location. *Angle Orthod* 47, 25-33 (1977).
- 59) Witt, E.: Möglichkeiten und Grenzen der kieferorthopädischen Behandlung Erwachsener. *Fortschr Kieferorthop* 52, 1-7 (1991)
- 60) Witt, E.: Behandlungskonzepte. In Miethke, R.R., D. Drescher (Hrsg.): *Kleines Lehrbuch der Angel-Klasse II,1 unter besonderer Berücksichtigung der Behandlung*. Quintessenz, Berlin (1996).