

Ausgabe: KN 9-2019, S. 12ff.

Thema: Kieferorthopädisch-kieferchirurgische Kombinationsbehandlung

Autoren: Dr. Ioan Barbur, Dr. Adina Barbur, Dr. Alexandra-Iulia Aghiorghiesei,
Dr. Alexandra Osan, Simion Bran und Dr. Mihaela Baciut.

Literatur

1. Yu D, Wang F, Wang X, Fang B, Shen SG. Presurgical motivations, self-esteem and oral health of orthognathic surgery patients. *The Journal of Craniofacial Surgery*. 2013;24(3):743-
2. Lu CH, Ko EW, Huang CS. The accuracy of video imaging prediction in soft tissue outcome after bimaxillary orthognathic surgery. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons*. 2003;61(3):333-42.
3. Van Hemelen G, Van Genechten M, Renier L, Desmedt M, Verbruggen E, Nadjmi N. Three-dimensional virtual planning in orthognathic surgery enhances the accuracy of soft tissue prediction. *Journal of cranio-maxillo-facial surgery: official publication of the European Association for Cranio-Maxillo-Facial Surgery*. 2015;43(6):918-25.
4. Solano-Hernández B, Antonarakis GS, Scolozzi P, Kiliaridis S. Combined Orthodontic and Orthognathic Surgical Treatment for the Correction of Skeletal Anterior Open-Bite Malocclusion: A Systematic Review on Vertical Stability. *J Oral Maxillofac Surg*. Januar 2013;71(1):98-109.
5. Jae Hyun Park, Michael Papademetriou, Carolyn Gardiner, John Grubbb. Anterior open bite correction with 2-jaw orthognathic surgery. *Am J Orthod Dentofacial Orthop*. Januar 2019;155(1):108-116.
6. Kuroda, S., Sugawara, Y., Tamamura, N., & Takano-Yamamoto, T. (2007). Anterior open bite with temporomandibular disorder treated with titanium screw anchorage: Evaluation of morphological and functional improvement. *Am J Orthod Dentofacial Orthop*. 2007 Apr; 131(4):550–560.
7. Yu D, Wang F, Wang X, Fang B, Shen SG. Presurgical motivations, self-esteem, and oral health of orthognathic surgery patients. *The Journal of craniofacial surgery*. 2013;24(3):743-8.
8. McGrath C, Pang HN, Lo ECM, King NM, Hagg U, Samman N. (2008). Translation and evaluation of a Chinese version of the Child Oral Health-related Quality of Life measure. *International Journal of Paediatric Dentistry*, 18(4), 267–274.
9. Storms AS, et al. Three-dimensional aesthetic assessment of class II patients before and after orthognathic surgery and its association with quantitative surgical changes, *Int J Oral Maxillofac Surg* (2017).

10. Park JH, Papademetriou M, Gardiner C, Grubb J. Anterior open bite correction with 2-jaw orthognathic surgery. *Am J Orthod Dentofacial Orthop*. Januar 2019;155(1):108-116
11. Kuroda S, Sakai Y, Tamamura N, Deguchi T, Takano-Yamamoto T. Treatment of severe anterior open bite with skeletal anchorage in adults: comparison with orthognathic surgery outcomes. *Am J Orthod Dentofacial Orthop* 2007; 132: 599-605.
12. Luther F, Morris DO, Hart C. Orthodontic preparation for orthognathic surgery: how long does it take and why? A retrospective study. *The British journal of oral & maxillofacial surgery*. 2003;41(6):401-6.
13. Finelle G. Digital Smile Design in interdisciplinary and orthodontic dental treatment planning. *J Dentofacial Anom Orthod* 2017;20:303.