

Ausgabe: face & body 2/2020

Thema: Anwendungsmöglichkeiten der Eigenfett-Therapie in der Gesichtschirurgie

Autor: S. S. Nagel, B. Thomas, D. Kotsougiani-Fischer, U. Kneser, C. Hirche

Literatur

1. Herold C, Ueberreiter K, Cromme F, Grimme M, Vogt PM. [Is there a need for intrapectoral injection in autologous fat transplantation to the breast? - An MRI volumetric study]. *Handchir Mikrochir Plast Chir.* 2011;43(2):119-124.
2. Eto H, Suga H, Matsumoto D, et al. Characterization of structure and cellular components of aspirated and excised adipose tissue. *Plastic and reconstructive surgery.* 2009;124(4):1087-1097.
3. Suga H, Matsumoto D, Inoue K, et al. Numerical measurement of viable and nonviable adipocytes and other cellular components in aspirated fat tissue. *Plastic and reconstructive surgery.* 2008;122(1):103-114.
4. To K, Crowley C, Lim SK, Khan WS. Autologous adipose tissue grafting for the management of the painful scar. *Cytherapy.* 2019;21(11):1151-1160.
5. Kilroy GE, Foster SJ, Wu X, et al. Cytokine profile of human adipose-derived stem cells: expression of angiogenic, hematopoietic, and pro-inflammatory factors. *Journal of cellular physiology.* 2007;212(3):702-709.
6. Rehman J, Traktuev D, Li J, et al. Secretion of angiogenic and antiapoptotic factors by human adipose stromal cells. *Circulation.* 2004;109(10):1292-1298.
7. Jiang T, Xie Y, Zhu M, et al. The second fat graft has significantly better outcome than the first fat graft for Romberg syndrome: A study of three-dimensional volumetric analysis. *J Plast Reconstr Aesthet Surg.* 2016;69(12):1621-1626.
8. Eto H, Suga H, Inoue K, et al. Adipose injury-associated factors mitigate hypoxia in ischemic tissues through activation of adipose-derived stem/progenitor/stromal cells and induction of angiogenesis. *The American journal of pathology.* 2011;178(5):2322-2332.
9. Eto H, Kato H, Suga H, et al. The fate of adipocytes after nonvascularized fat grafting: evidence of early death and replacement of adipocytes. *Plastic and reconstructive surgery.* 2012;129(5):1081-1092.
10. Denadai R, Raposo-Amaral CA, Pinho AS, Lameiro TM, Buzzo CL, Raposo-Amaral CE. Predictors of Autologous Free Fat Graft Retention

in the Management of Craniofacial Contour Deformities. *Plastic and reconstructive surgery*. 2017;140(1):50e-61e.

11. Mashiko T, Yoshimura K. How does fat survive and remodel after grafting? *Clin Plast Surg*. 2015;42(2):181-190.
12. https://www.awmf.org/uploads/tx_szleitlinien/009-0171_S2k_Autologe-Fetttransplantation_2016-07-verlaengert.pdf. 11/2015.
13. Tonnard P, Verpaele A, Peeters G, Hamdi M, Cornelissen M, Declercq H. Nanofat grafting: basic research and clinical applications. *Plastic and reconstructive surgery*. 2013;132(4):1017-1026.
14. Masia J, Bordoni D, Pons G, Liuzza C, Castagnetti F, Falco G. Oncological safety of breast cancer patients undergoing free-flap reconstruction and lipofilling. *European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology*. 2015;41(5):612-616.
15. Sanzenbacher R, Frech, M. Streitpunkt Eigenfettbehandlung: Wenn Gewebe zur Arznei wird. *Deutsches Ärzteblatt*. 2019;1-2/2019(116(1-2): A-28 / B-24 / C-24):28-30.
16. Prantl L, Giunta, R., Horch, R., Herold, C., von Hassel, J. Streitpunkt Eigenfettbehandlung: Gewebe und nicht Arzneimittel. *Deutsches Ärzteblatt*. 2019;1-2/2019(116(1-2): A-26 / B-22 / C-22):26-28.
17. Gesundheit Bf. Zweiter Bericht der Bundesregierung über die Situation der Versorgung der Bevölkerung mit Geweben und Gewebezubereitungen. 2018; <https://www.bundesgesundheitsministerium.de/service/begriffe-von-a-z/g/gewebeprodukte.html> Accessed 19.12.2018.
18. Deutsche Gesellschaft der Plastischen RuÄCD. Leitlinie Autologe Fetttransplantation. 2018.