

## LITERATUR

**Ausgabe:** Implantologie Journal 6/20

**Thema:** Die Anwendung von autogenem Dentin – Alveolenmanagement mit histologischer Aufbereitung

**Autoren:** Andreas van Orten, M.Sc., M.Sc., Dirk Krischik, Dr. med. dent., M.Sc., M.Sc., Prof. Dr. Dr. Werner Götz

---

1) Schropp L, Wenzel A, Kostopoulos L & Karring T. Bone healing and soft tissue contour changes following single-tooth extraction: a clinical and radiographic 12-month prospective study. *Int J Periodontics Restorative Dent* 2003; 23 : 313–323.

2) Araujo M G & Lindhe J. Dimensional ridge alterations following tooth extraction: an experimental study in the dog. *J Clin Periodontol* 2005; 32: 212–218.

3) Nevins M, Camelo M, De Paoli S, Friedland B, Schenk R K, Parma-Benfenati S, Simion M, Tinti C & Wagenberg B. A study of the fate of the buccal wall of extraction sockets of teeth with prominent roots. *Int J Periodontics Restorative Dent* 2006; 26: 19–29.

4) Fickl S, Zuhr O, Wachtel H, Stappert C F, Stein J M and Hürzeler M B. Dimensional changes of alveolar ridge contour after different socket preservation techniques. *J Clin Periodontol* 2008; 35: 906 – 913.

5) Allrad R H, Lekkas C and Swart J G. Autologous versus homologous bone grafting in osteotomies, secondary cleft repairs and ridge augmentations : a clinical study. *Oral Surg Oral Med Oral Pathol* 1987;64: 269-274.

6) Glass Y, Eickholz P, Nentwig H G, Dannewitz B, Glossar der Grundbegriffe für die Praxis. Knochenersatz- und aufbaumaterialien. *Parodontologie* 2008;19: 465-474.

7) Bartee B K. Extraction site reconstruction for alveolar ridge preservation. Part 1: rationale and materials selection. *J Oral Implantol* 2001;27: 187–193.

8) Pang K-M et al. Autogenous demineralized dentin matrix from extracted tooth fort he augmentation of alveolar bone defect: a prospective randomized clinical trial in comparison with anorganic bovine bone. *Clin Oral Impl Res* 2017;28: 809-815.

- 9) Cardaropoli D et al.: New bone formation using an extracted tooth as a biomaterial: A case report with histologic evidence. In *J Periodont Res* 2019;39: 157-163.
- 10) Kim Y K, Kim S G, Byeon J H, Lee H J, Um I U, Lim S C , & Kim S Y. Development of a novel bone grafting material using autogenous teeth. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, 2010; 109(4): 496–503.
- 11) Kim Y K et al. Tooth-derived bone graft material. *J Korean Assoc Oral Maxillofac Surg* 2013; 39: 103-111.
- 12) Um I W, Kim Y K, Park J C, & Lee, J H. Clinical application of autogenous demineralized dentin matrix loaded with recombinant human bone morphogenetic-2 for socket preservation: A case series. *Clin Implant Dent Relat Res* 2019; 21(1): 4-10.
- 13) Nampo T, Watahiki J, Enomoto A, Taguchi T, Ono M, Nakano H, Maki K. A new method for alveolar bone repair using extracted teeth for the graft material. *J Periodontol* 2010; 81: 1264-1272.
- 14) Bindermann I, Hallel G, Nardy C, Yaffe A, Sapoznikov L. A novel procedure to process extracted teeth for immediate grafting of autogenous dentin. *J Interdiscipl Med Dent Sci* 2014; 2: 1-5.
- 15) Schwarz F, Schmucker A, & Becker J. Initial case report of an extracted tooth root used for lateral alveolar ridge augmentation. *J Clin Periodontol* 2016; 43: 985-989.
- 16) van Orten A. Die Anwendung von A- und i-PRF in der zahnärztlichen Praxis – eine Standortbestimmung, *Dent Implant* 2018: 362-373.
- 17) Miron R J, Zucchelli G, Pikos M A, Salama M, Lee S, Guillemette V, Choukroun J. Use of platelet-rich fibrin in regenerative dentistry: a systematic review. *Clin Oral Investig* 2017; 21(6): 1913-1927.
- 18) Ghanaati S, Al-Maawi S. Autologe Alternative. *Teamwork* 2018; 3: 190-197.
- 19) Toloue S M, Chesnoiu-Matei I, Blanchard S B. A clinical and histomorphometric study of calcium sulfate compared with freeze-dried bone allograft for alveolar ridge preservation. *J Periodontol* 2012; 83: 847–855.
- 20) Cardaropoli G, Araujo M & Lindhe J. Dynamics of bone tissue formation in tooth extraction sites. An experimental study in dogs. *J Clin Periodontol* 2003; 30: 809–818.
- 21) Areewong K, Chantaramungkorn M, Khongkhunthian P. Platelet-rich fibrin to preserve alveolar bone sockets following tooth extraction: A randomized controlled trial. *Clin Implant Dent Relat Res* 2019: 1-8.

- 22) Buser D, Chen S T, Weber H P & Belser U C. Early implant placement following single- tooth extraction in the esthetic zone: biological rationale and surgical procedures. *Int J Periodontics Restorative Dent* 2008;28: 441–451.
- 23) Lekovic V, Kenney E B, Weinlaender M, Han T, Klokkevold P, Nedic M & Orsini M. A bone regenerative approach to alveolar ridge maintenance following tooth extraction. Report of 10 cases. *J Periodontol* 1997;68: 563–570.
- 24) van der Weijden F, Dell’acqua F & Slot D E. Alveolar bone dimensional changes of post-extraction sockets in humans: a systematic review. *J of Clin Periodontol* 2009; 36: 1048–1058.
- 25) Becker W, Hujoel P and Becker B E. Effect of barrier membranes and autologous bone grafts on ridge width preservation around implants. *Clin Implant Dent Relat Res* 2002; 4: 143-149.
- 26) Koga T et al.: Bone regeneration using dentin matrix depends on the degree of demineralization and particle size. *PLoS One* 2016: 1-12.
- 27) Schwarz F et al.: Zahnwurzeln als Augmentate. *ZWR* 1 2016; 106: 70-74.
- 28) Ramanauskaite A, Sahin D, Sader R, Becker J, & Schwarz F: Efficacy of autogenous teeth for the reconstruction of alveolar ridge deficiencies: a systematic review. *Clin Oral Investig* 2019; 23(12): 4263-4287.