

References

Er:YAG laser scanner

for implant site preparation

Drs Norberto Berna, Giovanni Olivi, Luca Marigo & Massimo Cordaro, Italy

implants – international magazine of oral implantology 3/20

1. Schwarz F. and Olivier W. Influence of implant bed preparation using an Er:YAG laser on the osseointegration of titanium implants: a histomorphometrical study in dogs Journal of Oral Rehabilitation 2007;4:273-281.
2. Stübinger S, von Rechenberg vB, Zeilhofer HF, Sader R, Landes C. Er:YAG laser osteotomy for removal of impacted teeth: clinical comparison of two techniques. Lasers in Surgery and Medicine 2007;39(7) 583-588.
3. Salina S, Maiorana C, Iezzi G , Colombo A, Fontana F, Piattelli A. Histological evaluation, in rabbit tibiae, of osseointegration of mini-implants in sites prepared with Er:YAG laser versus sites prepared with traditional burs. Journal of long-term effects of medical implants 2006;16(2):145-156.
4. Yoshino T, Aoki A, Oda S, et al. Long-term histologic analysis of bone tissue alteration and healing following Er:YAG laser irradiation compared to electrosurgery. Journal of Periodontology 2009;80(1) 82-92.
5. Stübinger S, Ghanaati S, Saldamli B, Kirkpatrick CJ, Sader R. Er:YAG laser osteotomy: preliminary clinical and histological results of a new technique for contact-free bone surgery. Eur Surg Res. 2009; 42(3):150-6.
6. Akyol UK, Güngörümüs M. Histologic evaluation of the effects of Er:YAG laser on bone ablation. J Contemp Dent Pract. 2009;10(5):65-72.
7. Maglione M, Bevilacqua et al observational study on the preparation of the implant site with Piezosurgery vs Drill: Comparison between the two methods in terms of postoperative pain, Surgical Time and operational Advantages.
8. Panduric DG, Juric IB, Music S, Molčanov K, Sušić M, Anic I. Morphological and ultrastructural comparative analysis of bone tissue after Er:YAG laser and surgical drill osteotomy. Photomed Laser Surg. 2014 Jul;32(7):401-8. doi: 10.1089/pho.2014.3711.
9. Blaskovic M, Gabrić D, Coleman NJ, Slipper IJ, , Gjorgjevska E. Bone Healing Following Different Types of Osteotomy: Scanning Electron Microscopy (SEM) and Three-Dimensional SEM Analyses. Microsc Microanal. 2016 Dec;22(6):1170-1178.
10. Kesler G, Romanos G, Koren R. Use of Er:YAG laser to improve osseointegration of titanium alloy implants-a comparison of bone healing. Int J Oral Maxillofac Implants. 2006;21(3):375-9

11. Augello M, Cattin C, et al. Comparative microstructural analysis of bone osteotomies after cutting by computer-assisted robot-guided laser osteotome and piezoelectric osteotome: an in vivo animal study. *Laser in medical scienze* 33(3) April 2018
12. Herrero-Climent M, Santos-García R, Jaramillo-Santos R, et al. Assessment of Osstell ISQ's reliability for implant stability measurement: a cross-sectional clinical study. *Med Oral Patol Oral Cir Bucal.* 2013;18(6):877–882.
13. Zix J, Kessler-Liechti G, Mericske-Stern R. Stability measurements of 1-stage implants in the maxilla by means of resonance frequency analysis: a pilot study. *Int J Oral Maxillofac Implants.* 2005;20:747–752.
14. Sim CP, Lang NP. Factors influencing resonance frequency analysis assessed by Osstell mentor during implant tissue integration: I. Instrument positioning, bone structure, implant length. *Clin Oral Implants Res.* 2010;21:598–604.
15. Trisi P, Carlesi T, Colagiovanni M, Perfetti G. Implant stability quotient (ISQ) vs direct in vitro measurement of primary stability (micromotion):effect of bone density and insertion torque. *J Osteol Biomat.* 2010;1:141–151.
16. Kesler G, Shvero DK, Tov YS, Romanos G. Platelet derived growth factor secretion and bone healing after Er:YAG laser bone irradiation. *J Oral Implantol.* 2011;37(3) 195-204.