

Ausgabe: cosmetic dentistry 4/2020
Thema: Korrektur eines skelettal offenen Bisses mit kombiniertem Therapiekonzept
Autoren: Prof. Dr. Nezar Watted, Prof. Dr. Mohamad Abu-Hussein Mohamad, Dr. Obeida Awad, Prof. Dr. Dr. Peter Proff, Dr. Dr. Ali Watted, Dr. Mohamad Abu Yunis

Literatur

- 1 Baek, S.H.; Ahn, H.W.; Kwon, Y.H. & Choi, J.Y.: Surgery First Approach in Skeletal Class III Malocclusion Treated with 2-Jaw Surgery: Evaluation of Surgical Movement and Postoperative Orthodontic Treatment. Journal of Craniofacial Surgery, 2010, .21: 332-338
- 2 Buddgaard M, Melsen B, Terp S. Changes during and following total maxillary osteotomy (Le Fort I Procedure): a cephalometric study. Eur J Orthod 1986; 8: 21-29.
- 3 Burstone GJ. The integumental profile. Am J Orthod 1958; 44: 1-25.
- 4 Burstone GJ. Lip posture and its significance in treatment planning. Am J Orthod 1967; 53: 262-84.
- 5 Calangna L, Silverman S, Garfinkel L. Influence of neuromuscular conditioning on centric relation registrations. J Prosthet Dent 1973; 30: 598-604.
- 6 Celenza, F V. The centric position. replacement and character. J Prosthet Dent 1973; 30: 591-598.
- 7 Coh-Stock C. Die chirurgische immediate Regulierung der Kiefer, chirurgische Behandlung der Prognathie. Dtsch Vjschr Zahnheik 1921; 37: 320-331.
- 8 Collins P, Epker BN. The alar basse Cinch: A technique for prevention of alar base flaring secondary to maxillary surgery. Oral Surg 1982; 53: 549-554.
- 9 Dal Pont G. L`osteotomia retromolare per la correzione della progenia, Minerva chir.,1959; 18: 1138-1141.
- 10 Dal Pont G. Die retromolare Osteotomie zur Korrektur der Progenie, der Retrogenie und des Mordex apertus. Öst Z Stoma 1961; 58: 8-10.
- 11 Epker BN, Fish LC. Surgical-orthodontic correction of open-bite deformity. Am J Orthod 1977; 71: 278-299.
- 12 Epker BN, Fish LC. Surgical superior repositioning of the maxilla: What to do with the mandibile? Am j Orthod 1980; 78: 164-191.
- 13 Friede H, Kahnberg K-E, Adell R, Riedell A. Accuracy of cephalometric prediction in orthognatic surgery. J Oral Maxillofac Surg 1987;45: 754-760.
- 14 Hogeman KE, Sarnäs KV, Willmar K. One-stage surgical and dental-orthopedic correction of bimaxillary and craniofacial dysostosis. In: Fortschr Kiefer Gesichtsch, Bd 18: 39-49, Thieme Verlag, Stuttgart, 1974.

15 Kondo E. & Aoba, T.J. (2000). Nonsurgical and Nonextraction Treatment of Skeletal Class III Open Bite: Its Long-Term Stability. *American Journal of Orthodontics and Dentofacial Orthopedics*, 2000, 117: 267-287

16 Kondo E. & Arai, S: Nonsurgical and Nonextraction Treatment of a Skeletal Class III Adult Patient with Severe Prognathic Mandible. *World Journal of Orthodontics*, 2005, .6: 233-247.

17 Lee DY, Bailey LJ, Proffit WR. Soft tissue changes after superior repositioning of the Maxilla with Le Fort I osteotomy: 5-years follow-up. *Int J Adult Orthod Orthognath* 1996; 11: 301-311.

18 Legan HL, Burstone GJ. Soft tissue cephalometric analysis for orthognathic surgery. *J Oral Surg* 1980; 38: 744-51.

19 O'Brien, K.; Wright, J.; Conboy, F.; Appelbe, P.; Bearn, D.; Caldwell, S.; Harrison, J.; Hussain, J.; Lewis, D.; Littlewood, S.; Mandall, N.; Morris, T.; Murray, A.; Oskouei, M.; Rudge, S.; Sandler, J.; Thiruvenkatachari, B.; Walsh, T. & Turbill, E.: Prospective, Multi-Center Study of the Effectiveness of Orthodontic/Orthognathic Surgery Care in the United Kingdom, *American Journal of Orthodontics and Dentofacial Orthopedics*, 2009, 135: 709-714.

20 Obwegeser H., Trauner R. Zur Operationstechnik bei der Progenie und anderen Unterkieferanomalien. *Dtsch Zahn Mund Kieferheilk* 1955; 23: 1-17.

21 Obwegeser H. The surgical correction of mandibular prognathism and retrognathia with consideration of genioplasty. *J Oral Surg* 1957; 10: 687-692.

22 Radney LJ, Jacobs JD. Soft tissue changes associated with surgical total maxillary intrusion. *Am J Orthodont* 1981; 80: 191-212.

23 Rekow ED, Speidel TM, Koenig RA. Location of the mandibular center of Autorotation in maxillary impaction surgery. *Am J Orthod Dentofac Orthop* 1993; 103: 530-536.

24 Rosen HM. Lip-nasal esthetics following Le Fort I Osteotomy. *Plast Reconstr Surg* 1988; 81: 171-182.

25 Schuchardt K. Formen des offenen Bisses und ihre operativen Behandlungsmöglichkeiten. *Fortsch Kiefer Gesichtschir. Bd 1*, Thieme Verlag, Stuttgart, 1955.

26 Stansbury CD, Evans CA, Miloro M, BeGole EA, Morris DE.: Stability of open bite correction with sagittal split osteotomy and closing rotation of the mandible. *J Oral Maxillofac Surg* 2010;68:149-159.

27 Watted N, Bill J. Untergesichtsverlängerung bei der Klasse-II-Dysgnathien: Neues von der kieferorthopädisch-kieferchirurgischen Therapie. *Ästhe Zahnmed* 1999; 4: 1992-300.

28 Watted N, Witt E, Bill J. Therapy Concept for the combined orthodontic-surgical Treatment of Angle Class II Deformities with Short-face Syndrome: New Aspects for surgical Lengthening of the Lower Face. *Clin Orthod research*, 2000; 3:78-93.

29 Watted N, Bill J, Witt E, Reuther J. Lengthening of the lower face Angle class II patients with skelettaly deep bite (short-face-syndrome) through combined orthodontic-surgical treatment. 75th Congress of the European Orthodontic Society Strasbuorg, France1999.

30 Williamsone EH, Evans DL, Barton WA, Williams BH. The effect of bite plane use on terminal hinge axis location, Angle Orthod. 1977; 47: 25-33.

31 Williamsone EH, Caves SA, Edenfield RJ, Morse PK. Cephalometric analysis: comparisons between maximum intercuspation and centric relation, Am J Orthod 1978; 74: 672-677.