

Ausgabe: Oralchirurgie Journal 1/21

Thema: Parodontitis. Lokalantibiotika: gezielte Wirkung ohne Belastung und Disziplin

Autor: Kulzer GmbH

Literatur

1. Repräsentative Umfrage unter Parodontitis-Betroffenen, Kantar GmbH, Januar 2020.
2. Kim TS et al., J Periodontol, 2002 Nov. 73 (11) 1285 – 91. Pharmacokinetic profile of a locally administered doxycycline gel on crevicular fluid, blood, and saliva.
3. Leyer et al. Adjunctive Topical Application of 14 % Doxycycline-Gel. An Observational Study. J Dent Res (Spec Iss 92A): 2186, 2013.
4. Kim TS, Bürklin T, Lee SH, Eickholz P, Zimmer H, Kim CK: Systemic detection of doxycycline after local administration Acta Odontologica Scandinavica, 2009, Vol. 67, No. 5, Pages 289 – 296.
5. Interne Daten, Kulzer GmbH.
6. Preshaw PM et al., J Clin Periodontal. 2004 Sept;31 (9): 697 – 707: Subantimicrobial dose doxycycline as adjunctive treatment for periodontitis. A review.
7. Metzger Z et al., Int Endod J. 2008 Apr; 41 (4): 303 – 9: Low-dose doxycycline inhibits bone resorption associated with apical periodontitis.