

**Ausgabe:** Endodontie Journal 1/2021

**Thema:** Komplexe Revision eines Unterkiefermolaren mit fünf Kanälen

**Autor:** Dr. Bernhard Bengs

---

### Literatur

- 1 Tahmasbi M, Jalali P, Nair MK, Barghan S, Nair UP. Prevalence of Middle Mesial Canals and Isthmi in the Mesial Root of Mandibular Molars: An In Vivo Cone-beam-Computed Tomographic Study. J Endod 2017 Jul; 43(7): 1080-1083.
- 2 Nosrat A, Deschenes RJ, Tordik PA, Hicks ML, Fouad AF. Middle mesial canals in mandibular molars: incidence and related factors. J Endod 2015 Jan; 41(1): 28-32.
- 3 Jang YE, Kim BS, Kim Y. Clinical Factors Associated with apical Periodontitis Visible on Cone-beam Computed Tomography but missed with Periapical Radiographs: A Retrospective Clinical Study. J Endod 2020 Jun; 46(6): 832-838.
- 4 Weinberg EM, Pereda AE, Khurana S, Lotlikar PP, Falcon C, Hirschberg C. Incidence of Middle Mesial Canals Based on Distance between Mesial Canal Orifices in Mandibular Molars: A Clinical and Cone-beam Computed Tomographic Analysis. J Endod 2020 Jan; 46(1): 40-43.
- 5 Akbarzadeh N, Aminoshariae A, Khalighinejad N, Palomo JM, Syed A, Kulild JC, Sadeghi G, Mickel A. The Association between the Anatomic Landmarks of the Pulp Chamber Floor and the Prevalence of Middle Mesial Canals in Mandibular First Molars: An In Vivo Analysis. J Endod 2017 Nov; 43(11): 1797-1801.
- 6 Azim AA, Deutsch AS, Solomon, CS. Prevalence of middle mesial canals in mandibular molars after guided troughing under high magnification: an in vivo investigation. J Endod 2015 Feb; 41(2): 164-168.