

LITERATUR

Ausgabe: Oralchirurgie Journal 2/21

Thema: Augmentation mit biphasischem, phykogenem KAM

Autor: Dr. med. dent. Alessandro Hellmuth Ponte

[1] Petrakakis P, Bergmann A, Langzeitbetrachtung zu Implantatüberlebensraten mit und ohne Augmentation, Praktische Implantologie und Implantatprothetik pip, 2017(1):64-67.

[2] Schopper C, Ziya-Ghazvini F, Goriwoda W, et al. HA/TCP compounding of a porous CaP biomaterial improves bone formation and slows degradation – a long-term histological study. J Biomed Mater Res B Appl Biomater 2005;74(1):458-67.

[3] Spassova E, Gintenreiter S, Halwax E, et al. Chemistry, ultrastructure and porosity of monophasic and biphasic bone forming materials derived from marine algae. Mat.-wiss. u Werksto tech. 2007;38(12):1-8.

[4] Zhou AJ, Clokie CM, Peel SA. Bone formation in algae-derived and synthetic calcium phosphates with or without poloxamer. J Craniofac Surg 2013;24(2):354-9.

[5] Zhou AJ, Peel SA, Clokie CM. An evaluation of hydroxyapatite and biphasic calcium phosphate in combination with Pluronic F127 and BMP on bone repair. J Craniofac Surg 2007;18(6):1264-75.

[6] Schopper C, Moser D, Spassova E, et al. Bone regeneration using a naturally grown HA/TCP carrier loaded with rhBMP-2 is independent of barrier-membrane effects. J Biomed Mater Res A 2008;85(4):954-63.

[7] Khoury F, Happe A. Soft tissue management in oral implantology: a review of surgical techniques for shaping an esthetic and functional peri-implant soft tissue structure. Quintessence Int Jul-Aug 2000;31(7):483-99.

[8] Puisys A, Linkevicius T. The influence of mucosal tissue thickening on crestal bone stability around bone-level implants. A prospective controlled clinical trial. *Clin Oral Implants Res* 2015;26:123-129.