

## Literature

### **Clinical relevance of the use of implant-supported provisional restorations to contour the emergence profile**

*Dr Marina Siegenthaler, Switzerland*

*implants – international magazine of oral implantology 3/22*

---

1. Chen, S. T., & Buser, D. (2014). Esthetic outcomes following immediate and early implant placement in the anterior maxilla--a systematic review. *Int J Oral Maxillofac Implants, 29 Suppl*, 186-215. doi:10.11607/jomi.2014suppl.g3.3
2. Small, P. N., & Tarnow, D. P. (2000). Gingival recession around implants: a 1-year longitudinal prospective study. *International Journal of Oral and Maxillofacial Implants, 15*(4), 527-532.
3. Castellon, P., Casadaban, M., & Block, M. S. (2005). Techniques to facilitate provisionalization of implant restorations. *Journal of Oral and Maxillofacial Surgery, 63*(9 Suppl 2), 72-79. doi:10.1016/j.joms.2005.05.152
4. Siegenthaler M, Strauss FJ, Gamper F, Hämmerle CH, Jung RE, Thoma DS. Anterior implant restorations with a convex emergence profile increase the frequency of recession: 12-month results of a randomized controlled clinical-trial. *J Clin Periodontol* (accepted 2022)
5. Chu, S. J. (2020). Inverted body-shift concept in macroimplant design to enhance biologic and esthetic outcomes: A clinical report. *Journal of Prosthetic Dentistry*. doi:10.1016/j.prosdent.2020.09.018
6. Steigmann, M., Monje, A., Chan, H. L., & Wang, H. L. (2014). Emergence profile design based on implant position in the esthetic zone. *International Journal of Periodontics and Restorative Dentistry, 34*(4), 559-563. doi:10.11607/prd.2016