

LITERATUR

Ausgabe: Implantologie Journal 3/24

Thema: Weitere Langzeitstudie: Keine Periimplantitis um zweiteiliges Implantatsystem

Autoren: Johannes Liebsch

1. Karapataki S, Vegh D, Payer M, Fahrenholz H, Antonoglou GN. Clinical performance of two-piece zirconia dental implants after 5 and up to 12 years. *Int J Oral Maxillofac Implants* 2023;38:1105–1114. DOI: 10.11607/jomi.10284
2. Brunello G, Rauch N, Becker K, Hakimi AR, Schwartz F, Becker J. Two-piece zirconia implants in posterior regions: A cohort study with a follow-up period of 9 years. *Clin Oral Implants Res* 2022;33:1233–1244.
3. Derks J, Tomasi C. Peri-implant health and disease. A systematic review of current epidemiology. *J Clin Periodontol* 2015;42(16):s158–s171.
4. Lee CT, Huang YW, Zhu L, Weltman R. Prevalences of peri-implantitis and peri-implant mucositis: Systematic review and meta-analysis. *J Dent* 2017;62:1–12.
5. Abdelhay N., Prasad S. & Gibson MP (2021) Failure rates associated with guided versus non-guided dental implant placement: a systematic review and meta-analysis. *BDJ Open* 7; 31 (2021). DOI: 10.1038/s41405-021-00086-1
6. Buser D, Weber HP, Lang NP. Tissue integration of non-submerged implants. 1-year results of a prospective study with 100 ITI hollow-cylinder and hollow-screw implants. *Clin Oral Implants Res* 1990;1:33–40.
7. Kohal RJ, Knauf M, Larsson B, Sahlin H, Butz F. One-piece zirconia oral implants: One-year results from a prospective cohort study. 1. Single tooth replacement. *J Clin Periodontol* 2012;39:590–597.
8. Greene JC, Vermillion JR. The simplified oral hygiene index. *J Am Dent Assoc* 1964;68:7–13.