

Ohrläppchenrekonstruktion bei Patienten mit 'Tunnel-Piercing'

Autoren: Dr. med. Daniel Sattler, Dr. med. Hans-Georg Damert, Dr. med. Silke Altmann, Dr. med. Armin Kraus, Prof. Dr. med. Manfred Infanger

Literatur

- [1] Williams AM, Majumder S. Earlobe reconstruction following plug and tunnel piercing. *J Plast Reconstr Aesthet Surg.* 2010 Mar;63(3):e294-5.
- [2] Agarwal R. Repair of cleft earlobe using double opposing Z-plasty. *Plast Reconstr Surg.* 1998 Oct;102(5):1759-60.
- [3] Chiummariello S, Iera M, Arleo S, Alfano C. L-specular plasty versus double-round plasty: two new techniques for earlobe split repair. *Aesthetic Plast Surg.* 2011 Jun;35(3):398-401.
- [4] Oh DY, Kim SW, Ahn ST, Rhie JW. Correction of earlobe cleft with tongue-in-groove technique. *J Craniofac Surg.* 2011 Sep;22(5):1785-7.
- [5] Yabe T, Muraoka M. Double opposing V-Y hinge flap. *Ann Plast Surg.* 2003 Dec;51(6):641-2.
- [6] Maral T, Tuncali D, Ozgur F, Gursu KG. A technique for the repair of simple congenital earlobe clefts. *Ann Plast Surg.* 1996 Sep;37(3):326-31.
- [7] Vujevich J, Goldberg LH, Obagi S. Repair of partial and complete earlobe clefts: a review of 21 methods. *J Drugs Dermatol.* 2007 Jul;6(7):695-9.
- [8] Zoltie N. Split earlobes: a method of repair preserving the hole. *Plast Reconstr Surg.* 1987 Oct;80(4):619-21.
- [9] Munro-Ashman D, MacDonald A, Feiweil M. Split earlobe syndrome. *Contact Dermatitis.* 1975 Dec;1(6):393-4.