

LITERATUR

Thema: Neues Kombi-Transplantat in Biomaterialien

Autor: CAMLOG Vertriebs GmbH

1. Cerabone® plus usability test.
2. 78.5% of users reported easier or much easier application compared to particulate material without hyaluronic acid; Data on file: Customer survey among 156 clinicians.
3. Tadic et al. Comparison of different methods for the preparation of porous bone substitution materials and structural investigations by synchrotron μ -computer tomography. Mat.-wiss. u. Werkstofftech. 2004, 35, No. 4.
4. Seidel and Dingeldein 2004. cerabone® – Bovine Based Spongiosa Ceramic Seidel et al. Mat.-wiss. u. Werkstofftech. 35:208–212.
5. Kyyak et al. Hyaluronic Acid with Bone Substitutes Enhance Angiogenesis In Vivo. Materials (Basel) 2022. 15(11):3839.
6. Kyyak et al. The Influence of Hyaluronic Acid Biofunctionalization of a Bovine Bone Substitute on Osteoblast Activity In Vitro. Materials (Basel). 2021. 14(11):2885.
7. Qasim SSB, Trajkovski B, Zafiropoulos GG. The response of human osteoblasts on bovine xenografts with and without hyaluronate used in bone augmentation. J Biomater Sci Polym Ed. 2024 Apr;35(6):880- 897. doi: 10.1080/09205063.2024.2311454. Epub 2024 Feb 12. PMID: 38346177.
8. Zhao, N., Wang, X., Qin, L., Zhai, M., Yuan, J., Chen, J., & Li, D. (2016). Effect of hyaluronic acid in bone formation and its applications in dentistry. Journal of biomedical materials research Part A, 104(6), 1560-1569.
9. Tawil et al. 2018. Sinus Floor Elevation Using the Lateral Approach and Window Repositioning and a Xenogeneic Bone Substitute as a Grafting Material: A Histologic, Histomorphometric, and Radiographic Analysis. Int J Oral Maxillofac Implants.33(5):1089-1096.

10. Riachi et al. 2012. Influence of material properties on rate of resorption of two bone graft materials after sinus lift using radiographic assessment. *Int J Dent.* 2012;737262.
11. Kloss et al. First Clinical Case Report of a Xenograft-Allograft Combination for Alveolar Ridge Augmentation Using a Bovine Bone Substitute Material with Hyaluronate (Cerabone® Plus) Combined with Allogeneic Bone Granules (Maxgraft®). *J Clin Med.* 2023. 12(19):6214.
12. Brown et al. New studies on the heat resistance of hamster-adapted scrapie agent: threshold survival after ashing at 600 degrees C suggests an inorganic template of replication, *PNAS* 2000. 97(7): 3418–3421.
13. Pröhl A et al. In Vivo Analysis of the Biocompatibility and Bone Healing Capacity of a Novel Bone Grafting Material Combined with Hyaluronic Acid. *Int J Mol Sci.* 2021. 22(9):48
14. Rakašević et al. Reconstructive Peri-Implantitis Therapy by Using Bovine Bone Substitute with or without Hyaluronic Acid: A Randomized Clinical Controlled Pilot Study. *J Funct Biomater.* 2023 Mar 8;14(3):149.
15. Rayahin, J. E., Buhrman, J. S., Zhang, Y., Koh, T. J., & Gemeinhart, R. A. (2015). High and low molecular weight hyaluronic acid differentially influence macrophage activation. *ACS biomaterials science & engineering*, 1(7), 481-493.
16. Pirnazar P. et al. 'Bacteriostatic effects of hyaluronic acid. *Journal of Periodontology* 1999. 70:370-374.
17. Håkansson et al. Regulation of granulocyte function by hyaluronic acid. In vitro and in vivo effects on phagocytosis, locomotion, and metabolism. *J Clin Invest.* 1980;66:298–305.
18. Wisniewski HG, Vilcek J. TSG-6: An IL-1/TNF-inducible protein with anti-inflammatory activity. *Cytokine Growth Factor Rev.* 1997. 8:143-56.
19. Larjava et al. Characterization of one phenotype of human periodontal granulation-tissue fibroblasts. *J Dent Res.* 1989. 68:20-25.
20. Bartold PM, Page RC. The effect of chronic inflammation on gingival connective tissue proteoglycans and hyaluronic acid. *J Oral Pathol.* 1986. 15:367-74.

21. Bertolami CN, Messadi DV. The role of proteoglycans in hard and soft tissue repair. *Crit Rev Oral Biol Med.* 1994. 5:311-37.
22. Ruggiero et al. Hyaluronidase activity of rabbit skin wound granulation tissue fibroblasts. *J Dent Res.* 1987. 66:1283-7.
23. Mendes et al. Sodium hyaluronate accelerates the healing process in tooth sockets of rats. *Arch Oral Biol.* 2008. 53:1155-62.