

Bisskraft, Stabilität und FKO-Apparaturen

Ein Beitrag von Dr. Gregory S. Antonarakis (Universität Genf/Schweiz), Prof. Dr. Heidrun Kjellberg (Universität Göteborg/Schweden) und Prof. Dr. Stavros Kiliaridis (Universität Genf/Schweiz).

Literatur

1. Bondevik O. Changes in occlusion between 23 and 34 years. *Angle Orthod* 1998;68:75-80
2. Ormiston JP, Huang GJ, Little RM, Decker JD, Seuk GD. Retrospective analysis of long-term stable and unstable orthodontic treatment outcomes. *Am J Orthod Dentofacial Orthop* 2006;128:568-574
3. Lerstøl M, Torgot O, Vandevska-Radunovic V. Long-term stability of dentoalveolar and skeletal changes after activator-headgear treatment. *Eur J Orthod* 2010;32:28-35
4. Melrose C, Millett DT. Toward a perspective on orthodontic retention. *Am J Orthod Dentofacial Orthop* 1998;113:507-514
5. Nanda R, Burzin J, Kim JM. Factors influencing the stability of Orthodontically treated dentitions. In: Hösl E, Baldauf A (eds). *Retention and long-term stability*. 8th International Conference for Orthodontists 1993; Hüthig, Heidelberg, pp. 41-46
6. Pancherz H. The nature of Class II relapse after Herbst appliance treatment: a cephalometric long-term investigation. *Am J Orthod Dentofacial Orthop* 1991;100:220-233
7. Wieslander L. Long-term effect of treatment with the headgear-Herbst appliance in the early mixed dentition. Stability or relapse? *Am J Orthod Dentofacial Orthop* 1993;104:319-329
8. Graber TM, Neumann B. The activator: use and modifications. In: Graber TM and Neumann B (eds) *Removable orthodontic appliances*. 1997; Saunders, Philadelphia, pp. 213-215
9. Helkimo E, Carlsson GE, Carmeli Y. Bite force in patients with functional disturbances of the masticatory system. *J Oral Rehabil* 1975;2:397-406
10. Staudt CB, Kiliaridis S. Association between mandibular asymmetry and occlusal asymmetry in young adult males with Class III malocclusion. *Acta Odontol Scand* 2010;68:131-140
11. Thilander B, Pena L, Infante C, Parada SS, de Mayorga C. Prevalence of malocclusion and orthodontic treatment need in children and adolescents in Bogota, Colombia. An epidemiological study related to different stages of dental development. *Eur J Orthod* 2001;23:153-167
12. Pancherz H, Anehus M. Masticatory function after activator treatment. An analysis of masticatory efficiency, occlusal conditions and EMG activity. *Acta Odontol Scand* 1978;36:309-316
13. Kiliaridis S, Mills CM, Antonarakis GS. Masseter muscle thickness as a predictive variable in treatment outcome of the twin-block appliance and masseteric thickness changes during treatment. *Orthod Craniofac Res* 2010;13:203-213
14. Antonarakis GS, Kjellberg H, Kiliaridis S. Predictive value of molar bite force on Class II functional appliance treatment outcomes. *Eur J Orthod* 2010;34:244-249
15. Björk A, Skjeller V. Facial development and tooth eruption. An implant study at the age of puberty. *Am J Orthod* 1972;62:339-383
16. Ingervall B, Helkimo E. Masticatory muscle force and facial morphology in man. *Arch Oral Biol* 1978;23:203-206