

Thema: Sofortimplantation in der ästhetischen Zone

Autoren: ZA Felix Knoch, Dr. Luís Bessa, Dr. Marcus Striegel

Literatur:

1. Chappuis V, Engel O, Reyes M, Shahim K, Nolte L-P, Buser D Ridge Alterations Postextraction in the Esthetic Zone: A 3D Analysis with CBCT CLINICAL INVESTIGATIONS. JDR Clin. Res. Suppl. 92.
2. Gomez-Meda R, Esquivel J, Blatz MB The esthetic biological contour concept for implant restoration emergence profile design. (2021).
3. Hsu Y-T, Lin G-H, Wang H-L Effects of Platform-Switching on Peri-implant Soft and Hard Tissue Outcomes: A Systematic Review and Meta-analysis. Int J Oral Maxillofac Implants 32:e9–e24 (2017).
4. Rosa J, Rosa A, Francischone C, Sotto-Maior B Diameter selection of implants placed in extraction sockets: A new approach. Dent Press Implantol 8:80–89 (2014).
5. Tavelli L, Barootchi S, Ravidà A, Oh TJ, Wang HL What Is the Safety Zone for Palatal Soft Tissue Graft Harvesting Based on the Locations of the Greater Palatine Artery and Foramen? A Systematic Review. J Oral Maxillofac Surg 77:271.e1-271.e9 (2019).
6. Yoshino S, Kan MSDJYK, Rungcharassaeng MSK, Roe P, Lozada MSJL Effects of Connective Tissue Grafting on the Facial Gingival Level Following Single Immediate Implant Placement and Provisionalization in the Esthetic Zone: A 1-Year Randomized Controlled Prospective Study. Int. J. Oral Maxillofac. Implants 29 (2014).