

Ausgabe: BDIZ konkret 2/2025

Titel: Crestales Hard- und Weichgewebsmanagement aktueller Implantatsysteme

Autoren: Prof. Jörg Neugebauer, Dres Steffen Kistler & Frank Kistler & Prof. Günter Dhom

Literatur

1. Agustin-Panadero R, Leon-Martinez R, Labaig-Rueda C et al. Influence of Implant-Prosthetic Connection on Peri-Implant Bone Loss: A Prospective Clinical Trial with 2-Year Follow-up. Int J Oral Maxillofac Implants 2019; 34: 963–968.
2. Berglundh T, Lindhe J. Dimension of the periimplant mucosa. Biological width revisited. J Clin Periodontol 1996; 23: 971–973.
3. Canullo L, Iurlaro G, Iannello G. Double-blind randomized controlled trial study on post-extraction immediately restored implants using the switching platform concept: soft tissue response. Preliminary report. Clin Oral Implants Res 2009; 20: 414–420.
4. Coelho PG, Jimbo R, Tovar N et al. Osseointegration: hierarchical designing encompassing the micrometer, millimeter, and nanometer length scales. Dent Mater 2015; 31: 37–52.
5. Hamilton A, Putra A, Nakapaksin P et al. Implant prosthodontic design as a predisposing or precipitating factor for peri-implant disease: A review. Clinical Implant Dentistry and Related Research 2023; 25: 710–722.
6. Hermann JS, Buser D, Schenk RK et al. Biologic Width around one- and two-piece titanium implants. Clin Oral Implants Res 2001; 12: 559–571.
7. Howe MS. Implant maintenance treatment and peri-implant health. Evid Based Dent 2017; 18: 8–10.
8. Kistler F, Frank I, Kistler S et al. Clinical Guidelines for Minimizing Complications in Subcrestal Implant Placement. In 40. Annual Meeting Academy of Osseointegration, Seattle, WA, USA: 2025.
9. Linkevicius T, Linkevicius R, Gineviciute E et al. The influence of new immediate tissue level abutment on crestal bone stability of subcrestally placed implants: A 1-year randomized controlled clinical trial. Clin Implant Dent Relat Res 2021; 23: 259–269.
10. Linkevicius T, Puisys A, Linkevicius R et al. The influence of submerged healing abutment or subcrestal implant placement on soft tissue thickness and crestal bone stability. A 2-year randomized clinical trial. Clin Implant Dent Relat Res 2020; 22: 497–506.

11. Lombardi T, Berton F, Salgarello S et al. Factors Influencing Early Marginal Bone Loss around Dental Implants Positioned Subcrestally: A Multicenter Prospective Clinical Study. *J Clin Med* 2019; 8.
12. Lombardo G, Corrocher G, Pighi J et al. The impact of subcrestal placement on short locking-taper implants placed in posterior maxilla and mandible: a retrospective evaluation on hard and soft tissues stability after 2 years of loading. *Minerva Stomatol* 2014; 63: 391–402.
13. Novaes Junior AB, de Oliveira RR, Taba Junior M et al. Crestal bone loss minimized when following the crestal preparation protocol: a histomorphometric study in dogs. *J Oral Implantol* 2005; 31: 276–282.
14. Pozzi A, Tallarico M, Moy PK. Three-year post-loading results of a randomised, controlled, split-mouth trial comparing implants with different prosthetic interfaces and design in partially posterior edentulous mandibles. *Eur J Oral Implantol* 2014; 7: 47–61.
15. Ramanauskaite A, Rocuzzo A, Schwarz F. A systematic review on the influence of the horizontal distance between two adjacent implants inserted in the anterior maxilla on the inter-implant mucosa fill. *Clin Oral Implants Res* 2018; 29 Suppl 15: 62–70.
16. Romanos GE, Delgado-Ruiz R, Sculean A. Concepts for prevention of complications in implant therapy. *Periodontol 2000* 2019; 81: 7–17.
17. Rothamel D, Heinz M, Ferrari D et al. Impact of machined versus structured implant shoulder designs on crestal bone level changes: a randomized, controlled, multicenter study. *Int J Implant Dent* 2022; 8: 31.
18. Rupp F, Liang L, Geis-Gerstorfer J et al. Surface characteristics of dental implants: A review. *Dent Mater* 2018; 34: 40–57.
19. Schoenbaum TR, Karateew ED, Schmidt A et al. Implant-Abutment Connections and Their Effect on Implant Survival Rates and Changes in Marginal Bone Levels (Delta): A Systematic Review and Meta-Analysis of 45,347 Oral Implants. *Int J Oral Maxillofac Implants* 2023; 38: 37–45.
20. Schwarz F, Hegewald A, Becker J. Impact of implant-abutment connection and positioning of the machined collar/microgap on crestal bone level changes: a systematic review. *Clin Oral Implants Res* 2014; 25: 417–425.
21. Schwarz F, Messias A, Sanz-Sanchez I et al. Influence of implant neck and abutment characteristics on peri-implant tissue health and stability. Oral reconstruction foundation consensus report. *Clin Oral Implants Res* 2019; 30: 588–593.
22. Tarnow DP, Cho SC, Wallace SS. The effect of inter-implant distance on the height of inter-implant bone crest. *J Periodontol* 2000; 71: 546–549.

23. Vervaeke S, Dierens M, Besseler J et al. The influence of initial soft tissue thickness on peri-implant bone remodeling. *Clin Implant Dent Relat Res* 2014; 16: 238–247.
24. Weigl P, Strangio A. The impact of immediately placed and restored single-tooth implants on hard and soft tissues in the anterior maxilla. *Eur J Oral Implantol* 2016; 9 Suppl 1: S89–106.
25. Weng D, Nagata MJ, Bell M et al. Influence of microgap location and configuration on the periimplant bone morphology in submerged implants. An experimental study in dogs. *Clin Oral Implants Res* 2008; 19: 1141–1147.
26. Weng D, Nagata MJ, Leite CM et al. Influence of microgap location and configuration on radiographic bone loss in nonsubmerged implants: an experimental study in dogs. *The International journal of prosthodontics* 2011; 24: 445–452.
27. Wennerberg A, Sennerby L, Kultje C et al. Some soft tissue characteristics at implant abutments with different surface topography. A study in humans. *J Clin Periodontol* 2003; 30: 88–94.
28. Yoo J, Moon IS, Yun JH et al. Effect of initial placement level and wall thickness on maintenance of the marginal bone level in implants with a conical implant-abutment interface: a 5-year retrospective study. *J Periodontal Implant Sci* 2019; 49: 185–192.