

Möglichkeiten und Grenzen der kieferorthopädischen Camouflage-Behandlung bei Patienten mit skelettaler Klasse III

von Prof. Dr. Bashar Muselmani

Literatur:

- 1- E. Steinerhäuser, I. Rudski-Janson, Kieferorthopädische Chirurgie , Band II, (286-292) quintessenz bibliothek 1994
- 2- William R. Proffit, Henry W. Fields, JR. David M. Sarver Contemporary Orthodontics, Fourth Edition: (640 -641) Mosby Elsevier. 2007
- 3- Moore T, Southard KA, Casco JS, et al. Buccal corridors and smile esthetics. Am J Orthod Dentofac Orthop 127:208-213, 2005.
- 4- Pancherz H, Fackel U. The skeletofacial growth pattern pre- and post-dentofacial orthopedics. Eur j Orthod 12:209-218, 1990.
- 5- Proffit WR, Phillips C, Douvartzidis N. *A comparison of outcomes of orthodontic and surgical-orthodontic treatment of Class III malocclusion in adults.* Am J Orthod Dentofacial Orthop. 1992;101(6):556-565.
- 6- Bailey LJ, Proffit WR, White RP. *Assessment of patients for orthognathic surgery.* Semin Orthod. 1999;5(4):209-222.
- 7- Ben-Bassat Y, Brin I. *Skeletal and dental patterns in patients with borderline Class III malocclusion.* Eur J Orthod. 2009;31: 327-333.
- 8- Lin J, Gu Y. *Lower incisor inclination and Holdaway angle as diagnostic keys in Class III treatment planning.* J Clin Orthod. 2003..
- 9- Tweed CH. *Indications for extraction in orthodontic therapy.* Am J Orthod. 1944.
- 10- Bell, W; B.Epker Surgical -orthodontic expansion of Maxilla. Am J Orthod 70, 517-518 (1979)

