

**Möglichkeiten und Grenzen der kieferorthopädischen Camouflage-Behandlung (Teil 2):
Behandlung von Patienten mit skelettalem offenem Biss**

von Prof. Dr. Bashar Muselmani

Literatur:

1. Proffit WR, Phillips C, Douvartzidis N. A comparison of outcomes of orthodontic and surgical-orthodontic treatment of Class III malocclusion in adults. *Am J Orthod Dentofacial Orthop.* 1992;101(6):556–565.
2. Proffit WR, Phillips C, Turvey TA. Stability after surgical-orthodontic correction of skeletal Class III malocclusion. *Angle Orthod.* 1991;61(1):31–41.
3. Bailey LJ, Proffit WR, White RP. Assessment of patients for orthognathic surgery. *Semin Orthod.* 1999;5(4):209–222.
4. Rakosi T. *Funktionelle Therapie in der Kieferorthopädie.* München; Wien: Hanser; 1985. S. 47–51, 431–442.
5. Kerr WJS, Miller S, Dawber JE. Class III malocclusion: surgery or orthodontics? *Br J Orthod.* 1992;19:21–24.
6. Ben-Bassat Y, Brin I. Skeletal and dental patterns in patients with borderline Class III malocclusion. *Eur J Orthod.* 2009;31(3):327–333.
7. Lin J, Gu Y. Diagnostic criteria for Class III malocclusion: Holdaway angle and incisor inclination. *J Clin Orthod.* 2003;37(10):589–595.
8. Janson G, Cattaneo PM, Henriques JF, de Freitas MR. Stability of Class III malocclusion treated with extraction and non-extraction protocols. *Am J Orthod Dentofacial Orthop.* 2004;126(4):476–482.
9. Graf H. *Rezidivprophylaxe bei kieferorthopädischer Therapie mit abnehmbaren Geräten.* Leipzig: Barth; 1979.
10. Kuroda S, Sugawara Y, Tamamura N, Takano-Yamamoto T. Treatment of severe anterior open bite with skeletal anchorage. *Am J Orthod Dentofacial Orthop.* 2007;131(4 Suppl):S61–S69.
11. Sugawara J, Baik UB, Umemori M, Takahashi I, Kawamura H, Mitani H. Treatment and posttreatment dentoalveolar changes following molar intrusion

with skeletal anchorage. *Am J Orthod Dentofacial Orthop.* 2002;121(6):566–575.

12. Sherwood KH, Burch JG, Thompson WJ. Closing anterior open bites by intruding molars with titanium miniplate anchorage. *Am J Orthod Dentofacial Orthop.* 2002;122(6):593–600.

13. Baek SH, Choi JY, Lee JH. Orthodontic treatment of skeletal open bite using skeletal anchorage. *Korean J Orthod.* 2010;40(2):79–90.

14. Kuroda S, Sugawara Y, Deguchi T, Kyung HM, Takano-Yamamoto T. Clinical use of miniscrew implants in orthodontics: open bite correction. *Angle Orthod.* 2007;77(6):1114–1120.

15. Gomes AF, et al. Orthodontic camouflage versus orthognathic surgery in adult Class III patients: systematic review. *Eur J Orthod.* 2020;42(5):531–540.

16. Reichenbach E, Mainhold G. Funktionelle Entwicklung des Schluckmusters im frühen Kindesalter. *Fortschr Kieferorthop.* 1970;31.

17. Schwarz AM. *Lehrbuch der Kieferorthopädie.* Leipzig: Thieme; 1956.

18. Damon DH. The Damon low friction bracket: a biologically compatible straight-wire system. *J Clin Orthod.* 1988;32:670–680.

19. Kuroda S, et al. Treatment of severe anterior open bite with skeletal anchorage in adults. *Am J Orthod Dentofacial Orthop.* 2007;132(5):599–605.

20. Melsen B. *Adult Orthodontics.* Oxford: Wiley-Blackwell; 2012.

21. Kim SH, et al. Skeletal open bite and its treatment stability. *Semin Orthod.* 2002;8(3):162–172.