

Aesthetics Follows Function – Extraktionstherapie auch aus ästhetischer Sicht?

von Dr. Eva Marie Lessmann und Prof. Dr. Angelika Stellzig-Eisenhauer.

Literaturverzeichnis

- [1] Angle, E. H.: Treatment of Malocclusion of the Teeth: Angle's System. 7. Aufl. S. S. White, Philadelphia (1907).
- [2] Boley, J. C., Pontier, J. P., Smith, S., Fulbright, M.: Facial changes in extraction and nonextraction patients. Angle Orthod 68:539–546 (1998).
- [3] Bowman, S. J., Johnston, L. E. Jr.: The esthetic impact of extraction and nonextraction treatments on Caucasian patients. Angle Orthod 70:3–10 (2000).
- [4] Elias, K. G., Sivamurthy, G., Bearn, D. R.: Extraction vs nonextraction orthodontic treatment: A systematic review and meta-analysis. Angle Orthod 94:83–106 (2024).
- [5] Iared, W., Koga da Silva, E. M., Iared, W., Rufino Macedo, C.: Esthetic perception of changes in facial profile resulting from orthodontic treatment with extraction of premolars: A systematic review. J Am Dent Assoc 148:9–16 (2017).
- [6] Kochar, G. D., Londhe, S., Chopra, S. S., Kohli, S., Kohli, V. S., Kamboj, A., Verma, M.: Treatment effects and lip profile changes following premolars extraction treatment vs fixed functional treatment in Class II division 1 malocclusion: A randomized controlled clinical trial. Dental Press J Orthod 28 (2023).
- [7] Peck, S.: Extractions, retention and stability: the search for orthodontic truth. Eur J Orthod 39:109–115 (2017).
- [8] Tweed, C. H.: Indications for the extraction of teeth in orthodontic procedure. Am J Orthod Oral Surg 30:405–428 (1944).