Titel: From urgent care to comprehensive oral rehabilitation
Interdisciplinary management of a patient with dental anxiety

Author: Lucas Fahling, Germany

References

[1] Wölber J., Frick K.; Motivierende Gesprächsführung in der zahnärztlichen Therapie; Zahnmedizin up2date 3; 2014,247–269

[2] Ramseier CA, Warnakulasuriya S, Needleman IG et al. Consensus Report: 2nd European Workshop on Tobacco Use Prevention and Cessation for Oral Health Professionals. Int Dent J 2010; 60: 3–6

[3] Hagenow F. Zahnbehandlungsangst: Evaluation eines Trainingskurses für Zahnärzte zum Umgang mit ängstlichen Patienten. Saarbrücken: Akademikerverlag; 2012

[4] Ridder P, Craniomandibuläre Dysfunktion: Interdisziplinäre Diagnose- und Behandlungsstrategien. München Urban und Fischer 2011

[5] Bernhöft K, Klammt J. Untersuchungen über Beziehungen zwischen funktionellen Störungen an den Kiefergelenken und der Halswirbelsäule. Zahn-, Mund- und Kieferheilkunde mit Zentralblatt; 1998; 76:36-9

[6] Losert-Bruggner B., Dudek B., Hülse M. Die craniomandibuläre Dysfunktion (CMD) Manuelle Medizin 2010, 5.2010.48: 343-352

[7] Ruge S., Quooß, A., Kordaß, B. (2010) Virtuelle Artikulation – Visualisierung dynamischer Okklusionsbeziehungen. Aus Lebendige Wissenschaft - Spitzenforschung in der Zahnheilkunde 2010:84–90

[8] Khoury F, Hanser T. Mandibular bone block harvesting from the retromolar region: a 10-year prospective clinical study. Int J Oral Maxillofac Implants. 2015; 30(3):688 697.

[9] Nissan J, Mardinger O, Calderon S, Romanos GE, Chaushu G. Cancellous bone block allografts for the augmentation of the anterior atrophic maxilla. Clin Implant Dent Relat Res 2011; 13(2):104-111.

[10] Khoury F., Khoury CH. Mandibular bone block grafts: diagnosis, instrumentation, harvesting techniques and surgical procedures. In: Khoury F, Antoun, H, Missika P, editors. Bone Augmentation in Oral Implantology. Berlin: Quintessence, 2007.

[11] Cordaro L, Torsello F, Accorsi Ribeiro C, Liberatore M, Mirisola di Torresanto V. Inlay-onlay grafting for three-dimensional reconstruction of the posterior atrophic maxilla with mandibular bone. Int J Oral Maxillofac Surg 2010;39:350-357.

[12] Turri A, Elgalil I, Vazirisani F, Johansson A, Emanuelsson L, Dahlin C, Thomsen P, Omar O. Guided bone regeneration is promoted by the molecular events in the membrane compartment. Biomaterials 84;2016: 167-183

[13] Nevins M, Mellonig JT, Clem DS, 3rd, Reiser GM, Buser DA. Implants in regenerated bone: long-term survival. Int J Periodontics Restorative Dent 1998;18:34-45.

[14] Peroz I, Imhoff B; S2k-Leitlinie Okklusionsschiene zur Behandlung craniomandibulärer Dysfunktionen und zur präprothetischen Therapie. (2024) AWMF Reg.-Nr. 083-051

[15] Plaster U. Natürliche Asymmetrien und die patientenindividuelle Wiedergabe der Okklusionsebene ohne traditionellen Transferbogen. Quintessenz Zahntech 2013;39:1266-1280

[16] Araujo NS, Moda MD, Silva EA, Zavanelli AC, Mazaro JV, Pellizer EP. Survival of all-ceramic restorations after a minimum follow-up of five years: a systematic review. Quintessence Int 2016;47:395-405.