OEMUS MEDIA AG 2026

Präzision statt Kompromiss – Die VILA als Therapiekonzept für komplexe kieferorthopädische Behandlungsfälle

von Dr. Franziska Lang[*], Dr. Norbert Lang[*], Dr. Antonia Böddeker[*], Priv.-Doz. Dr. Isabel Knaup[*], Prof. Dr. Dr. Ali Modabber[**], Prof. Dr. Dr. Frank Hölzle[**], Priv.-Doz. Dr. Rogerio B. Craveiro[*], Prof. Dr. Michael Wolf[*].

[*]Klinik für Kieferorthopädie, Universitätsklinik RWTH Aachen, Deutschland

[**]Klinik für Mund-, Kiefer- und Geschichtschirurgie, Universitätsklinik RWTH Aachen, Deutschland

Literatur:

[1] Grauer D, Proffit WR. Accuracy in tooth positioning with a fully customized lingual orthodontic appliance. American Journal of Orthodontics and Dentofacial Orthopedics 2011; 140: 433–443. doi:https://doi.org/10.1016/j.ajodo.2011.01.020

[2] Pauls A, Nienkemper M, Schwestka-Polly R, et al. Therapeutic accuracy of the completely customized lingual appliance WIN. Journal of Orofacial Orthopedics / Fortschritte der Kieferorthopädie 2017; 78: 52–61. doi:10.1007/s00056-016-0058-9

[3] Coenen FA, Bartz JR, Niederau C, et al. Orthodontic treatment quality evaluated by partially automated digital IOTN and PAR index determination: a retrospective multicentre study. Eur J Orthod 2023; 45: 308–316. doi:10.1093/ejo/cjac064

[4] Lang FA, Lang NA, Vorloeper J, et al. Validation of a digital, partly automated three-dimensional cast analysis for evaluation of orthodontic treatment assessment. Head Face Med 2025; 21. doi:10.1186/s13005-025-00515-8

[5] Gallone M, Robiony M, Bordonali D, et al. Multidisciplinary treatment with a customized lingual appliance for an adult patient with severe Class III malocclusion and multiple missing teeth. American Journal of Orthodontics and Dentofacial Orthopedics 2019; 156: 401–411. doi:10.1016/j.ajodo.2019.05.004

[6] Lang FA, Lang NA, Lode GI, et al. Digital evaluation with ABO eCRE of treatment quality and efficiency with completely customized lingual and prefabricated vestibular appliances in combined orthodontic-orthognathic therapy: a retrospective cohort study. Prog Orthod 2026; 27. doi:10.1186/s40510-026-00639-w

[7] Möhlhenrich SC, Steur J, Chhatwani S, et al. Occlusal changes during orthognathic therapy using a completely customised lingual appliance - initial retrospective observations. Clin Oral Investig 2025; 29. doi:10.1007/s00784-025-06347-9

[8] Klein KP, Kaban LB, Masoud MI. Orthognathic Surgery and Orthodontics: Inadequate Planning Leading to Complications or Unfavorable Results. Oral Maxillofac Surg Clin North Am 2020; 32: 71–82. doi:10.1016/J.COMS.2019.08.008

[9] Hamada Y, Timothius CJC, Shin D, et al. Canine impaction – A review of the prevalence, etiology, diagnosis and treatment. Semin Orthod 2019; 25: 117–123. doi:10.1053/j.sodo.2019.05.002

[10] Cernochova P, Cernoch C, Klimo Kanovska K, et al. Treatment options for impacted maxillary canines and occurrence of ankylotic and resorptive processes: a 20-year retrospective study. BMC Oral Health 2024; 24. doi:10.1186/s12903-024-04662-3

[11] Incerti-Parenti S, Checchi V, Ippolito DR, et al. Periodontal status after surgical-orthodontic treatment of labially impacted canines with different surgical techniques: A systematic review. American Journal of Orthodontics and Dentofacial Orthopedics 2016; 149: 463–472

[12] Rodríguez-Cárdenas YA, Arriola-Guillén LE, Aliaga-Del Castillo A, et al. Three-dimensional changes in root angulation of buccal versus palatal maxillary impacted canines after orthodontic traction: A retrospective before and after study. Int Orthod 2021; 19: 216–227. doi:10.1016/j.ortho.2021.04.003

[13] Schmid JQ, Bettenhäuser-Hartung L, Kanemeier M, et al. The incidence of gingival recession with non-surgical crossbite correction using completely customized lingual appliances versus surgically assisted rapid palatal expansion in adults: a cohort study. Prog Orthod 2025; 26. doi:10.1186/s40510-025-00568-0

[14] Janssens Y, Siekmann H, Canal P, et al. Quality of occlusal outcome in adult patients with posterior crossbite treated with completely customized lingual appliances and CAD/CAM archwires for maxillary expansion and mandibular compression compared to adult Class I patients: A retrospective study. Int Orthod 2025; 23. doi:10.1016/j.ortho.2025.101040

[15] Housley JA, Nanda RS, Currier GF, et al. Stability of transverse expansion in the mandibular arch. American Journal of Orthodontics and Dentofacial Orthopedics 2003; 124: 288–293. doi:10.1016/S0889-5406(03)00450-5

[16] Schmid JQ, Gerberding E, Hohoff A, et al. Non-Surgical Transversal Dentoalveolar Compensation with Completely Customized Lingual Appliances versus Surgically Assisted Rapid Palatal Expansion in Adults—The Amount of Posterior Crossbite Correction. J Pers Med 2022; 12. doi:10.3390/jpm12111893

[17] Schmid JQ, Gerberding E, Hohoff A, et al. Non-Surgical Transversal Dentoalveolar Compensation with Completely Customized Lingual Appliances versus Surgically Assisted Rapid Palatal Expansion in Adults—Tipping or Translation in Posterior Crossbite Correction? J Pers Med 2023; 13. doi:10.3390/jpm13050807

[18] Janssens Y, Foley PF, Beyling F, et al. Quality of occlusal outcome in adult Class II patients treated with completely customized lingual appliances and Class II elastics compared to adult Class I patients. Eur J Orthod 2024; 46. doi:10.1093/ejo/cjae031

[19] Wiechmann D, Schwestka-Polly R, Hohoff A. Herbst appliance in lingual orthodontics. American Journal of Orthodontics and Dentofacial Orthopedics 2008; 134: 439–446. doi:10.1016/j.ajodo.2007.09.015

[20] Alouini O, Knösel M, Blanck-Lubarsch M, et al. Controlling incisor torque with completely customized lingual appliances. Journal of Orofacial Orthopedics 2020; 81: 328–339. doi:10.1007/s00056-020-00231-9

[21] Thiem LC, Rank P, Schmid JQ, et al. Favourable dentoalveolar changes after lower premolar extractions for Class III camouflage with completely customized lingual appliances. Head Face Med 2024; 20. doi:10.1186/s13005-024-00459-5

[22] Lossdörfer S, Schwestka-Polly R, Wiechmann D. Inklinationskontrolle der unteren Frontzähne bei der konservativen Therapie der Klasse III mit einer vollständig individuellen lingualen Apparatur. Journal of Orofacial Orthopedics 2013; 74: 381–396. doi:10.1007/s00056-013-0163-y

[23] Wiechmann D, Leven R, Rank P, et al. Dentoalveolar process remodelling in the anterior mandible after Class III camouflage treatment with lower premolar extractions. Head Face Med 2025; 21. doi:10.1186/s13005-025-00493-x