

DT GER – AU – CH 7+8/2013, p. 19

Artikel:

Alarmierend: Trotz Desinfektion bis zu 70 Prozent Keimbelastung

Quellen:

¹ Compendium of Continuing Education in Dentistry, Compendium Supplement, April 2013, Volume 34, Issue

² Studienreport Universität Witten/Herdecke: Mikrobiologische Belastungen von Serviettenketten, Februar 2012

³ Molinari JA. Microbial contamination of patient napkin holders. The Dental Advisor. 2010;29:1-3.

⁴ Study: Bacteria found on 70% of dental bib holders [news release]. San Francisco, CA: DrBicuspid.com; March 19, 2012.
<http://www.drBicuspid.com/index.aspx?sec=sup&sub=hyg&pag=dis&ItemID=310087>.
Accessed February 19, 2013.

⁵ Study finds bib chain potential source of bacteria. Dental Health Magazine; March 16, 2010.
<http://worldental.org/dental-news/study-finds-bib-chain-potential-source-of-bacteria/1184>. Accessed February 19, 2013.

⁶ Alt-Holland A, Srinivasan S, Lucier R, et al. Do bib clips pose a cross-contamination risk at the dental clinic? Compend Contin Educ Dent. 2012; 33(9 suppl):1-7.

⁷ The complete genome sequence of *Propionibacterium acnes*, a commensal of human skin. Science 305(5684):671-673(2004).

⁸ Dentist's office a 'perfect storm' for HIV, hepatitis exposure, health official says, CNN, March 30, 2013.
<http://edition.cnn.com/2013/03/29/health/oklahoma-dental-warning>.

⁹ 60 Oklahoma Dental Patients Test Positive for Hepatitis, HIV, abc News, April 18, 2013.
<http://abcnews.go.com/Health/60-oklahoma-dental-patients-test-positive-hepatitishiv/story?id=18991527#.UXo3zMoX7sB>.