

## Literaturverzeichnis

# Die kieferorthopädische Behandlung erwachsener Patienten – eine interdisziplinäre Therapie

Ein Beitrag von Prof. Dr. med. dent. N. Watted, Priv.-Doz. Dr. med. Dr. med. dent. J. Bill, Dr. med. dent. S. Kleimann, Dr. med. dent. B. Shlomi und Dr. med. dent. V. Reiser.

- [1] Albino, J.E., Tedesco, L.: Esthetic need for orthodontic treatment. In Melsen B, editor: Current controversies in orthodontics. Chicago, Quintessence Publishing, pp. 11–24 (1994).
- [2] Bass, N.M.: Dento-facial orthopaedics in der correction of the skeletal II malocclusion. Br J Orthod 9, 3–8 (1982).
- [3] Berscheid, E., Gangestade, S: The social psychological implications of facial physical attractiveness. Clin Plast Surg 9, 289–296 (1982).
- [4] Canut, J.: Eine Analyse der dentofazialen Ästhetik. Inf Orthod Kieferorthop 28, 83–105 (1996).
- [5] Dryland-Vig, K.W.L., Ellis III, E.: Diagnosis and treatment planning for the surgical-orthodontic Patient. Cli Plast Surg 16, 645–658 (1989).
- [6] Farkas, L.G., Kolar, J.C.: Anthropometry and art in the aesthetics of women's face. Clin Plast Surg 14, 599–615 (1987).
- [7] Flanary, C.M., Barnwell, G.M., Alexander, J.M.: Patient perceptions of orthognathic surgery. Am J Orthod 88, 137–145 (1985).
- [8] Helm, S., Siersbaek-Nielsen, S., Skieller, V., Björk, A.: Skelatal maturation of the hand in relation to maximum puberal growth in body height. Danish Dental Jornal 75, 1223-1234 (1971).
- [9] Jacobson, A.: The influence of children's dentofacial appearance on their social attractiveness as judged by peers and lay adults. Am J Orthod 79, 399–415 (1981).
- [10] Kiyak, H.A., Hohl ,T., West, R.A.: Psychologie changes in orthognathic surgery patients: a 24-month follow-up. J Oral Maxillofac Surg 42, 506–512 (1984).
- [11] Legan, H.L., Burstone, G.J.: Soft tissue cephalometric analysis for orthognathic surgery. J Oral Surg 38, 744–51 (1980).
- [12] McNamara, J.A., McDougall, Jr.P.D., Dierks, J.M.: Arch with development in Class II patients treated with extraoral force and functional jaw orthodontics. Am J Orthodont 52, 353–359 (1966).

- [13] Petrovic, A.G., Stutzmann, J.: Reaktionsfähigkeit des tierischen und menschlichen Kondylenknorpels auf Zell- und Molekularebene im Lichte einer kybernetischen Auffassung des faszialen Wachstums. Fortschr Kieferorthop 49, 405–425 (1988).
- [14] Schwarz, A.M.: Die Röntgendiagnostik. Urban & Schwarzenberg, Wien (1958).
- [15] Scott, O., Kijak, H.A.: Treatment expectation versus outcomes among orthodontic surgery patients. Int J Adult Orthod Orthognath Surg 6, 247–255 (1991).
- [16] Watted, N., Witt, E.: NMR study of TNJ changes following functional orthopaedic treatment using the „Würzburg approach“, European Orthodontic Society (EOS) 74th Congress (1998).
- [17] Watted, N.: Behandlung von Klasse II-Dysgnathien – Funktionskieferorthopädische Therapie unter besonderer Berücksichtigung der dentofazialen Ästhetik, Kieferorthop 13, 193–208 (1999).
- [18] Watted, N., Bill, J., Witt, E.: Therapy Concept for the Combined Orthodontic-Surgical Treatment of Angle Class II Deformities with Short Face Syndrome New Aspects for Surgical Lengthening of the Lower Face. Clin. Orthod. Res. 3, 78–93 (2000).
- [19] Watted, N., Bill, J., Witt, E., Reuther, J.: Lengthening of the lower face Angle class II patients with skeletally deep bite (shortface-syndrome) through combined orthodontic-surgical treatment. 75th Congress of the European Orthodontic Society Strasbourg, France(1999).
- [20] Watted, N., Teuscher, T., Wieber, M.: Vertikaler Gesichtsaufbau und Planung kieferorthopädisch-kieferchirurgischer Kombinationsbehandlungen unter besonderer Berücksichtigung der dentofazialen Ästhetik. Kieferorthop 16, 29–44 (2000).
- [21] Witt, E.: Möglichkeiten und Grenzen der kieferorthopädischen Behandlung Erwachsener. Fortschr Kieferorthop 52, 1–7 (1991)
- [22] Witt, E.: Behandlungskonzepte. In Miethke, R.R., D. Drescher (Hrsg.): Kleines Lehrbuch der Angle-Klasse II,1 unter besonderer Berücksichtigung der Behandlung. Quintessenz, Berlin (1996).