

Ausgabe: Dental Tribune GER 1/2017

Thema: Bisphosphonate und andere antiresorptive Medikamente. Relevante Aspekte für den zahnärztlichen Praxisalltag

Autor: Priv.-Doz. Dr. Dr. Heinz-Theo Lübbbers, Winterthur, Schweiz

Literatur

- [1] Grötz KA, Piesold J-U, Al-Nawas B: Bisphosphonat-assoziierte Kiefernekrose (BP-ONJ) und andere Medikamenten-assoziierte Kiefernekrosen. AWMF-Leitlinien 2012;
- [2] Pfammatter C, Kühl S, Filippi A, Lambrecht JT: Patienten unter Bisphosphonattherapie. Teil 1: Bisphosphonate – chemische Formel, Wirkungsmechanismus, Nebenwirkungen und Risikofaktoren. Quintessenz 2011; 62(6): 795-801.
- [3] Marx RE, Sawatari Y, Fortin M, Broumand V: Bisphosphonate-induced exposed bone (osteonecrosis/osteopetrosis) of the jaws: risk factors, recognition, prevention, and treatment. Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons 2005; 63(11): 1567-1575.
- [4] Svejda B, Muschitz C, Gruber R, Brandtner C, Svejda C, Gasser RW, Santler G, Dimai HP: [Position paper on medication-related osteonecrosis of the jaw (MRONJ)]. Wien Med Wochenschr 2016; 166(1-2): 68-74.
- [5] Khan AA, Morrison A, Hanley DA, Felsenberg D, McCauley LK, O'Ryan F, Reid IR, Ruggiero SL, Taguchi A, Tetradiis S, Watts NB, Brandi ML, Peters E, Guise T, Eastell R, Cheung AM, Morin SN, Masri B, Cooper C, Morgan SL, Obermayer-Pietsch B, Langdahl BL, Al Dabagh R, Davison KS, Kendler DL, Sandor GK, Josse RG, Bhandari M, El Rabbany M, Pierroz DD, Sulimani R, Saunders DP, Brown JP, Compston J, International Task Force on Osteonecrosis of the J: Diagnosis and management of osteonecrosis of the jaw: a systematic review and international consensus. J Bone Miner Res 2015; 30(1): 3-23.
- [6] Ruggiero SL, Dodson TB, Fantasia J, Gooday R, Aghaloo T, Mehrotra B, O'Ryan F, American Association of O, Maxillofacial S: American Association of Oral and Maxillofacial Surgeons position paper on medication-related osteonecrosis of the jaw--2014 update. Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons 2014; 72(10): 1938-1956.
- [7] Walter C, Al-Nawas B, Frickhofen N, Gamm H, Beck J, Reinsch L, Blum C, Grotz KA, Wagner W: Prevalence of bisphosphonate associated osteonecrosis of the jaws in multiple myeloma patients. Head & face medicine 2010; 6(11).
- [8] Metzler P, Zemann W, Lübbbers H-T, Guggenberger R, Lüssi A, Obwegeser JA, Grätz KW, Jacobsen C: Bone mineral density measurements performed by cone-beam computed tomography in the bisphosphonate-related osteonecrosis-affected jaw. Oral Radiology 2012; 28(2): 101-108.
- [9] Grötz KA. ASORS-Laufzettel: Überweisung / Konsil vor antiresorptiver Therapie des Knochens (Bisphosphonate, Denosumab). 2012.
- [10] Röntgen WC: Ueber eine neue Art von Strahlen. (Vorläufige Mittheilung.). Sitzungsberichte der Physik-med Gesellschaft zu Würzburg 1895; 132-141.
- [11] Hounsfield GN: Computerized transverse axial scanning (tomography). 1. Description of system. Br J Radiol 1973; 46(552): 1016-1022.
- [12] Lubbers HT, Terzic A, Zemann W, Jacobsen C, Obwegeser J, Kruse A: Computer assisted maxillofacial surgery. Minerva chirurgica 2011; 66(5): 469-481.
- [13] Mozzo P, Procacci C, Tacconi A, Martini PT, Andreis IA: A new volumetric CT machine for dental imaging based on the cone-beam technique: preliminary results. Eur Radiol 1998; 8(9): 1558-1564.

- [14] Ludlow JB, Ivanovic M: Comparative dosimetry of dental CBCT devices and 64-slice CT for oral and maxillofacial radiology. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008; 106(1): 106-114.
- [15] Lubbers HT, Matthews F, Damerau G, Kruse AL, Obwegeser JA, Gratz KW, Eyrich GK: Anatomy of impacted lower third molars evaluated by computerized tomography: is there an indication for 3-dimensional imaging? *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics* 2011; 111(5): 547-550.
- [16] Eyrich G, Seifert B, Matthews F, Matthiessen U, Heusser CK, Kruse AL, Obwegeser JA, Lubbers HT: 3-Dimensional imaging for lower third molars: is there an implication for surgical removal? *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons* 2011; 69(7): 1867-1872.
- [17] Dula K, Benic GI, Bornstein M, Dagassan-Berndt D, Filippi A, Hicklin S, Kissling-Jeger F, Luebbers HT, Sculean A, Sequeira-Byron P, Walter C, Zehnder M: SADMFR Guidelines for the Use of Cone-Beam Computed Tomography/Digital Volume Tomography. *Swiss Dent J* 2015; 125(9): 945-953.
- [18] Dula K, Bornstein MM, Buser D, Dagassan-Berndt D, Ettlin DA, Filippi A, Gabiou F, Katsaros C, Krastl G, Lambrecht JT, Lauber R, Luebbers HT, Pazera P, Turp JC, Sadmfr: SADMFR guidelines for the use of Cone-Beam Computed Tomography/Digital Volume Tomography. *Swiss Dent J* 2014; 124(11): 1169-1183.
- [19] Fehm T, Beck V, Banys M, Lipp HP, Hairass M, Reinert S, Solomayer EF, Wallwiener D, Krimmel M: Bisphosphonate-induced osteonecrosis of the jaw (ONJ): Incidence and risk factors in patients with breast cancer and gynecological malignancies. *Gynecol Oncol* 2009; 112(3): 605-609.
- [20] Saad F, Brown JE, Van Poznak C, Ibrahim T, Stemmer SM, Stopeck AT, Diel IJ, Takahashi S, Shore N, Henry DH, Barrios CH, Facon T, Senecal F, Fizazi K, Zhou L, Daniels A, Carriere P, Dansey R: Incidence, risk factors, and outcomes of osteonecrosis of the jaw: integrated analysis from three blinded active-controlled phase III trials in cancer patients with bone metastases. *Ann Oncol* 2012; 23(5): 1341-1347.
- [21] Vahtsevanos K, Kyrgidis A, Verrou E, Katodritou E, Triaridis S, Andreadis CG, Boukovinas I, Koloutsos GE, Teleioudis Z, Kitikidou K, Paraskevopoulos P, Zervas K, Antoniades K: Longitudinal cohort study of risk factors in cancer patients of bisphosphonate-related osteonecrosis of the jaw. *J Clin Oncol* 2009; 27(32): 5356-5362.
- [22] Kunchur R, Need A, Hughes T, Goss A: Clinical investigation of C-terminal cross-linking telopeptide test in prevention and management of bisphosphonate-associated osteonecrosis of the jaws. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons* 2009; 67(6): 1167-1173.
- [23] Yamazaki T, Yamori M, Ishizaki T, Asai K, Goto K, Takahashi K, Nakayama T, Bessho K: Increased incidence of osteonecrosis of the jaw after tooth extraction in patients treated with bisphosphonates: a cohort study. *Int J Oral Maxillofac Surg* 2012; 41(11): 1397-1403.
- [24] Mozzati M, Arata V, Gallesio G: Tooth extraction in patients on zoledronic acid therapy. *Oral Oncol* 2012; 48(9): 817-821.
- [25] Scoletta M, Arata V, Arduino PG, Lerda E, Chieccchio A, Gallesio G, Scully C, Mozzati M: Tooth extractions in intravenous bisphosphonate-treated patients: a refined protocol. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons* 2013; 71(6): 994-999.
- [26] Grotz KA, Al-Nawas B, Terheyden H: Implantate und Bisphosphonat-Therapie. *Implantologie* 2013; 21(1): 53-59.
- [27] Grotz KA, Schmidt BLJ, Walter C, Al-Nawas B: In which bisphosphonate patients am I allowed to place implants? A systematic review. *Z Zahnärztl Impl* 2010; 26(2): 153-161.
- [28] Lubbers HT, Bergau T, Hoff J, von Jackowski J: Dokumentationspflicht in der Zahncärztpraxis. *ZAHN PRAX* 2012; 15(4): 260-265.

- [29] Khan AA, Rios LP, Sandor GK, Khan N, Peters E, Rahman MO, Clokie CM, Dore E, Dubois S: Bisphosphonate-associated osteonecrosis of the jaw in Ontario: a survey of oral and maxillofacial surgeons. *J Rheumatol* 2011; 38(7): 1396-1402.
- [30] Stockmann P, Vairaktaris E, Wehrhan F, Seiss M, Schwarz S, Spriewald B, Neukam FW, Nkenke E: Osteotomy and primary wound closure in bisphosphonate-associated osteonecrosis of the jaw: a prospective clinical study with 12 months follow-up. *Support Care Cancer* 2010; 18(4): 449-460.
- [31] Wutzl A, Pohl S, Sulzbacher I, Seemann R, Lauer G, Ewers R, Drach J, Klug C: Factors influencing surgical treatment of bisphosphonate-related osteonecrosis of the jaws. *Head & neck* 2012; 34(2): 194-200.
- [32] Barasch A, Cunha-Cruz J, Curro FA, Hujoel P, Sung AH, Vena D, Voinea-Griffin AE, Group CC, Beadnell S, Craig RG, DeRouen T, Desaranayake A, Gilbert A, Gilbert GH, Goldberg K, Hauley R, Hashimoto M, Holmes J, Latzke B, Leroux B, Lindblad A, Richman J, Safford M, Ship J, Thompson VP, Williams OD, Yin W: Risk factors for osteonecrosis of the jaws: a case-control study from the CONDOR dental PBRN. *J Dent Res* 2011; 90(4): 439-444.